

P.L. Mechanism of training-induced mitochondrial biogenesis: A possible involvement of reactive oxygen species

Hideki MATOBA

Faculty of Socio-Arts and Sciences, The University of Tokushima, Japan

Training-induced adaptation of the skeletal muscle was rigorously studied in mid-1960' through 1970'. Gollnick PD, Holloszy J and Saltin B were the most influential exercise physiologists. Investigations in these days revealed that the most important physiological feature of endurance training-induced adaptation was an increased oxidative enzyme activity of the skeletal muscle. The increased oxidative enzyme activity is a reflection of augmented mitochondrial biogenesis. From the 1990' and thereafter, research efforts have been directed to elucidate mechanisms of the training-induced mitochondrial biogenesis. However, the mechanism has not yet been fully elucidated.

Recently, Gomez-Cabrera et al. (2008) published their research which indicated that reactive oxygen species (ROS) produced during endurance training played a critical role for training-induced mitochondrial biogenesis. However, later studies including ours reported divergent results.

Further studies are definitely needed to determine if ROS is involved in the endurance training-induced mitochondrial biogenesis.

S.L. Molecular mechanisms of unloading-mediated muscle atrophy and development of its countermeasures

Takeshi NIKAWA

Department of Nutritional Physiology, Institute of Health Biosciences, The University of Tokushima Graduate School

Skeletal muscle atrophy caused by unloading is characterized by both decreased responsiveness to myogenic growth factors (e.g. IGF-1, insulin) and increased proteolysis. Here we show that unloading stress led to skeletal muscle atrophy through the induction and activation of the ubiquitin ligase Cbl-b. Upon induction, Cbl-b interacted with and degraded the IGF-1 signaling intermediate IRS-1. In turn, loss of IRS-1 activated the FOXO3-dependent induction of atrogen-1, a dominant mediator of proteolysis in atrophic muscle. Forced Cbl-b expression was sufficient to induce IRS-1 ubiquitination and atrophy in rat tibialis anterior muscle. Cbl-b-deficient mice were resistant to unloading-induced atrophy and loss of muscle function. Furthermore, a penta-peptide mimetic of tyrosine⁶⁰⁸-phosphorylated IRS-1 inhibited Cbl-b-mediated IRS-1 ubiquitination and strongly decreased Cbl-b-mediated induction of atrogen-1. Our results indicate that the Cbl-b-dependent destruction of IRS-1 is a critical dual mediator of both the growth refractory and proteolytic arms of the atrophic response. Inhibition of Cbl-b-mediated ubiquitination may be a new therapeutic strategy for unloading-mediated muscle atrophy.