

Jpn. J. Ent., 65(1): 168–170. March 25, 1997

On the Nest of *Stenopsocus externus* BANKS (Psocoptera: Stenopsocidae)^{1), 2)}

Kazunori YOSHIZAWA³⁾ and Toyohei SAIGUSA

Biosystematics Laboratory, Graduate School of Social
and Cultural Studies, Kyushu University,
Ropponmatsu, Fukuoka, 810 Japan

Key words: *Stenopsocus externus*; Stenopsocidae; Psocoptera; nest; behavior.

In this paper, we report the nest of *Stenopsocus externus* BANKS from Guizhou, China.

Some taxonomic groups of the Psocoptera are known to weave silk nests or webs. As NEW (1987) mentioned, this behavior is widely spread in various psocid groups, for example *Caecilius* (Caeciliusidae), *Pseudocaecilius* (Pseudocaeciliidae), *Ectopsocus* (Ectopsocidae), *Reuterella* (Elipsocidae) and *Psoculus* (Mesopsocidae). In the genus *Stenopsocus*, JENTSCH (1938) reported that *S. stigmatics* and *S. immaculatus* often stay under a web together.


On June 30, 1995, one of us, SAIGUSA, found many silk nests of a peculiar shape (Fig. 1–3) on upper surface of leaves of two groups of a bamboo species in Jiangkou, Guizhou, China. The nest consists of a densely woven flat whitish film, roughly circular or oval in shape, about 5–8 mm in diameter and with its margin radially expanded at several portions, with which the film is attached to the surface of a leaf (Fig. 1–3). Usually, one to several nymphs of a psocid were found on the leaf under the nest (Fig. 4), and a few adults were also observed there often together with nymphs. Judging from the material component of the nest, it cannot be assumed that the nest was woven by spiders, lepidopterous larvae, etc. Additionally, some adults were observed under very roughly woven web (Fig. 5), which seems a nest of the species under construction. Therefore we consider that the nest was woven by the psocid. This psocid was determined as *Stenopsocus externus* BANKS, 1937 by YOSHIZAWA. This species is widely distributed in southern China and Taiwan.

The present stenopsocid's nest seems peculiar in shape because the nests of *Reuterella helvimacula* (Elipsocidae) (YOSHIZAWA, 1996), *Haplophallus* spp. and *Aaroniella* spp. (Philotarsidae) (YOSHIZAWA's personal observation) and *Stenopsocus*

¹⁾ Contribution from the Biosystematics Laboratory, Graduate School of Social and Cultural Studies, Kyushu University, Fukuoka (No. 12).

²⁾ This study was partly supported by Grant-in-aid for International Scientific Research (Field Research), Ministry of Educations, Science, Sports & Culture, Japanese Government. (No. 07041141)

³⁾ Research Fellow of the Japan Society for the Promotion of Science.



Figs. 1–5. — 1, Nests (dried specimens). 2, Ditto, right one magnified. 3, A nest in the field. 4, Ditto, upper half of the nest removed. Two larvae were staying under the nest. 5, An adult stays under the nest under construction. (Fig. 3–5 were taken by SONY Handycam.)

stigmatics and *S. immaculatus* (Stenopsocidae) (JENTSCH, 1938) are rough and irregular in shape. On the other hand, the nest of *Pseudocaecilius citricola* (Pseudocaeciliidae) is very similar to the present stenopsocid's nest (TURNER and MOCKFORD, pers. comm.). NEW (1987) mentioned that webbing habit has independently evolved in a number of lineages of the Psocoptera and the shapes of nests are variable among the taxa. In the future, when phylogenetic relationships of the Psocoptera are understood and nest of many groups of psocids are studied, evolution of the nesting behavior of the Psocoptera will be discussed.

As SAIGUSA observed the psocids only for an hour, we cannot confirm the role of the nest. But, as mentioned by TURNER (1973), the nest may serve for a refuge against predators.

We thank to Dr. B. D. TURNER (King's College, University of London) and Dr. E. L. MOCKFORD (Illinois State University) for giving us valuable information on the nest of *Pseudocaecilius citricola*, and to Prof. T. NAITO (Faculty of Agriculture, Kobe University), a representative of the field research project in China, for his help during the survey. YOSHIZAWA thanks to Prof. H. SHIMA and Assoc. Prof. O. YATA

(Biosystematics Laboratory, Kyushu University) for their constant guidance and encouragement.

References

- JENTSCH, S., 1938. Beiträge zur Kenntnis der Überordnung Psocoidea. 3. Zur Copeognathenf-auna Nordwestfalens. *Abh. westfal. Prov. Mus. naturk.*, **9**(4): 3–42.
- NEW, T. R., 1987. Biology of the Psocoptera. *Orient. Ins.*, **21**: 1–109.
- TURNER, B. D., 1973. The distribution of webs and egg-batches of *Pseudocaecilius citricola* (ASHMEAD) (Insecta: Psocoptera: Pseudocaeciliidae) on the leaves of *Magnifera indica* L. and *Cordia collococca* L. in Jamaica, W. I. *J. nat. Hist.*, **7**: 561–565.
- YOSHIZAWA, K., 1996. First record of *Reuterella helvimacula* (Psocoptera: Elipsocidae) from Japan. *Jpn. J. Ent.*, **64**: 187–188.

(Received July 11, 1996; Accepted September 4, 1996)