

Kontyû, 38 (3): 205-221. 1970

NOTES ON THE CHRYSOMELIDAE FROM TAIWAN IV*†

By Shinsaku Kimoto

*Biological Laboratory, Pre-Medical Course, School of Medicine,
Kurume University, Kurume*

Subfamily Alticinae, I

Nonarthra cyaneum Baly, 1874

=*Nonarthra nigricolle* (Weise, 1889)

Taipingshan (=Taiheizan), Ilan Hsien (1 ex., 23. iii. 1933, K. Satô leg.).

Sungan (=Nihonmatsu) —Peikeng (=Hokukô), Miaoli Hsien (1 ex., 10. iv. 1967, T. Shirôzu leg.).

Tienhsiang, Hualien Hsien (1 ex., 18. vii. 1966, B.-S. Chang leg.).

Sungkang, Nantou Hsien (1 ex., 19. v. 1965, 1 ex., 31. v. 1965, 1 ex., 1. vi. 1965, T. Shirôzu leg.; 1 ex., 16. vii. 1966, H. Kamiya leg.).

Tsifeng, Nantou Hsien (2 exs., 10. vii. 1966, H. Kamiya leg.).

Taataka Pass (=Taataka-anbu), Chiayi Hsien (1 ex., 3. iv. 1967, T. Shirôzu leg.).

Distribution: Japan, Taiwan, China, Indo-China.

Nonarthra variabile Baly, 1862

=*Nonarthra formosense* Chûjô, 1935, Trans. Nat. Hist. Soc. Formosa 25 : 360

(Taiwan: Musha, Rimogan, Ehoshiyama). **New Synonymy**

Yangmingshan, Taipei Hsien (1 ex., 25. v. 1965, S. Asahina leg.; 1 ex., 24. vi. 1965, S. Kimoto leg.; 1 ex., 24. viii. 1966, H. Kamiya leg.).

Wulai, Taipei Hsien (1 ex., 27. v. 1965, K. Morimoto leg.; 1 ex., 25. viii. 1966, H. Kamiya leg.).

Ilan, Ilan Hsien (2 exs., 15. iv. 1965, Y. Hirashima leg.; 1 ex., 19. viii. 1966, H. Kamiya leg.).

Taruling—Tzeen, Hualien Hsien (2 exs., 17. vii. 1966, B.-S. Chang leg.).

Nanshanchi, Nantou Hsien (2 exs., 28. iv. 1965, 5 exs., 6. v. 1965, 1 ex., 17. vi. 1965, T. Shirôzu leg.; 2 exs., 21. v. 1965, B.-S. Chang leg.; 1 ex., 30. vi. 1965, S. Kimoto leg.).

Juisui (=Mizuho), Hualien Hsien (5 exs., 21-22. iv. 1932, J. L. Gressitt leg.).

Meichi, Nantou Hsien (1 ex., 12. v. 1965, 1 ex., 16. v. 1965, T. Shirôzu leg.).

* Partly supported by a grant from "Japan Society for the Promotion of Science" in connection with the Japan-U. S. Cooperative Science Program: Zoogeography and Ecology of Pacific Area Insects.

† Part 1, 1967, Kontyû 35 (4): 368-374; Part 2, 1969, Esakia 7: 1-68; Part 3, 1970, Kontyû 38 (2): 176-183.

- Penpuchi, Nantou Hsien (1 ex., 2. v. 1965, T. Shirôzu leg.).
 Wushe—Yushih, Nantou Hsien (1 ex., 3. v. 1965, T. Shirôzu leg.).
 Sungkang, Nantou Hsien (1 ex., 31. v. 1965, 2 exs., 1. vi. 1965, T. Shirôzu leg.).
 Fenchihu, Chiayi Hsien (1 ex., 9. vii. 1965, Y. Kurosawa leg.; 1 ex., 24. vii. 1966, H. Kamiya leg.).
 Kuantzuling, Tainan Hsien (1 ex., 7. iv. 1965, S. Miyamoto leg.).
 Kuanshan (=Riran), Taitung Hsien (1 ex., 19. iv. 1932, J. L. Gressitt leg.).
 Chulu, Taitung Hsien (4 exs., 9. viii. 1966, H. Kamiya leg.).
 Kueitzuchiaio (=Kurarur), Pingtung Hsien (1 ex., 1. ix. 1921, T. Esaki leg.).
 Distribution: India, Burma, Indo-China, S. China, Taiwan, Ryukyu Is.

***Nonarthra bimaculatum* n. sp.**

Head, pronotum and scutellum shining black, elytra shining black with bluish luster, and with a pair of reddish markings on middle of elytra; in most specimens pronotum with anterior angles dark reddish brown; antennae blackish brown with three basal joints paler; ventral surface of thorax entirely black, abdomen entirely yellowish brown; legs mostly black, with apices of femora and base of tibiae, together with entire tarsi more or less brownish.

Head finely and remotely punctate, with a broad, ill-defined, transversal excavation just above frontal tubercles, surface of frontal tubercles smooth, shining, separated to each other by a short but distinct longitudinal furrow at middle, frons finely and remotely punctate. Antennae robust, depressed, somewhat shorter than half length of body, first joint robust, long; second short, about half as long as first and much slender; third nearly $1\frac{1}{3}$ times as long as second and much robust; fourth nearly as long as third but slightly robust; fourth to eighth nearly as long as in length to each other but gradually widened as go to terminal joints; ninth nearly $1\frac{1}{2}$ times as long as eighth and bluntly pointed at apex. Prothorax distinctly broader than long, anterior margin generally rounded posteriorly but almost straight in median portion; lateral margins slightly rounded, posterior margin widely rounded posteriorly; anterior margin widely rounded posteriorly; anterior angles nearly rectangular, basal angles rounded; surface convex, very finely but remotely punctate. Scutellum trigonate, surface convex, with a few very fine punctures. Elytra oblong-oval, surface convex, rather strongly and closely punctate, punctures consisted by two sorts of distinct and fine ones.




Fig. 1. a, Antenna of *Nonarthra bimaculatum* n. sp.; b, Male genitalia of *Chaetocnema (Tlanoma) yiei* n. sp.

Length: 3.8 mm.

Holotype: Sungkang, Nantou Hsien (3. vi. 1965, B.-S. Chang leg.).

Paratypes: Taipingshan, Ilan Hsien (1 ex., 5. viii. 1967, B.-S. Chang leg.). Tabu (=Taiko) —Sungan (=Nihonmatsu), Miaoli Hsien (1 ex., 3. vi. 1965, T. Shirôzu leg.). Tungpu—Tzuchung, Lulinshan, Nantou Hsien (1 ex., 10. iv. 1965, T. Saigusa leg.).

This new species closely resembles *N. cyaneum* by its general characteristics and the coloration, but differs in having the antennae much slender and with a pair of reddish markings on the middle of elytra.

Distribution: Taiwan.

***Psylliodes difficilis* Baly, 1874**

=*Psylliodes gigantea* Chûjô, 1935, Trans. Nat. Hist. Soc. Formosa 25: 367

(Taiwan: Taitô, Kwarenkô). **New Synonymy**

Fenchihu, Chiayi Hsien (1 ex., 7. vii. 1965, 1 ex., 8. vii. 1965, S. Kimoto leg.).

According to the study of the type of *P. gigantea* Chûjô, preserved in the Taiwan Agricultural Research Institute, Taipei, this species is identical with *P. difficilis* Baly, occurring in Japan, China and some other areas. Also, there is no distinct difference on the male genitalia between the Japanese and the Taiwan populations.

Distribution: China, Japan, Ryukyu Is., Taiwan, Bonin Is., Volcano Is.

***Psylliodes angusticollis* Baly, 1874**

=*Psylliodes formosana* Chûjô, 1963, Ann. Hist.-Nat. Mus. Nat. Hungarici 55: 401 (Taiwan: Kosempo). **New Synonymy**

Yangmingshan, Taipei Hsien (2 exs., 26. vi. 1965, S. Kimoto leg.).

Wulai, Taipei Hsien (1 ex., 27. v. 1965, K. Morimoto leg.).

Sungkang, Nantou Hsien (1 ex., 29. vi. 1965, T. Yamasaki leg.).

Sungkang—Tsifeng, Nantou Hsien (9 exs., 29. vi. 1965, S. Kimoto leg.).

Yungfeng—Kunyang, Nantou Hsien (1 ex., 1. vii. 1965, S. Kimoto leg.).

Fenchihu, Chiayi Hsien (1 ex., 10. iv. 1965, S. Miyamoto leg.; 1 ex., 12. iv. 1965, T. Saigusa leg.; 14 exs., 7. vii. 1965, 1 ex., 8. vii. 1965, S. Kimoto leg.; 2 exs., 7. vii. 1965, T. Nakane leg.; 1 ex., 25. vii. 1966, H. Kamiya leg.).

Chiaoliping, Chiayi Hsien (1 ex., 13. iv. 1965, T. Saigusa leg.).

Alishan, Chiayi Hsien (3 exs., 6. vii. 1965, S. Kimoto leg.).

Through the kindness of Prof. M. Chûjô, Takamatsu, I had an opportunity to study the type of *P. formosana* Chûjô, preserved in his personal collection. The name given as *formosana* is no doubt a synonymy of *P. angusticollis* Baly. The type of *angusticollis* Baly is preserved in Brit. Mus. (Nat. Hist.).

Distribution: Japan, Ryukyu Is., Taiwan, Korea, Sachalin.

***Chaetocnema (Chaetocnema) formosensis* Chûjô, 1935**

Yangmingshan, Taipei Hsien (7 exs., 24. vi. 1965, S. Kimoto leg.).

Wulai, Taipei Hsien (1 ex., 22. viii. 1966, H. Kamiya leg.).

Kuanhsi, Hsinchu Hsien (1 ex., 16. iv. 1967, B.-S. Chang leg.).

Sungkang—Tsifeng, Nantou Hsien (1 ex., 29. vi. 1965, T. Nakane leg.).

Distribution: Taiwan, Ryukyu Is.

Chaetocnema (Tlanoma) basalis Baly, 1877

Yangmingshan, Taipei Hsien (1 ex., 24. vi. 1965, S. Kimoto leg.).

Tahu (=Taiko)—Sungan (=Nihonmatsu), Miaoli Hsien (1 ex., 9. iv. 1967, T. Shirôzu leg.).

Fenchihu, Chiayi Hsien (1 ex., 10. iv. 1965, T. Shirôzu leg.).

5-10km S. of Fenchihu, Chiayi Hsien (1 ex., 11. iv. 1965, Y. Hirashima leg.).

Distribution: India, Ceylon, Burma, Indo-China, S. China, Taiwan, Ryukyu Is.

Chaetocnema (Tlanoma) discreta Baly, 1877

Shuangchi, Hsintien, Taipei Hsien (2 exs., 19. vi. 1965, K. Kurosawa leg.).

Kuantzuling, Tainan Hsien (2 exs., 7. iv. 1965, S. Miyamoto leg.).

Distribution: Indo-China, China, Taiwan, Ryukyu Is., Japan.

Chaetocnema (Tlanoma) subcostata Jacoby, 1889

Chiaoliping, Chiayi Hsien (1 ex., 13. iv. 1965, T. Saigusa leg.).

Distribution: Burma, Taiwan.

Chaetocnema (Tlanoma) tonkinensis Chen, 1934

=*Chaetocnema (Tlanoma) tonkinensis taiwanensis* Chûjô, 1965, Spec. Bull. Lep. Soc. Japan. 1: 99 (Taiwan: Musha). **New Synonymy**

Yangmingshan, Taipei Hsien (1 ex., 28. iii. 1965, S. Miyamoto leg.; 1 ex., 25. v. 1965, Y. Miyatake leg.).

Ilan, Ilan Hsien (1 ex., 16. iv. 1965, Y. Hirashima leg.).

Distribution: Indo-China, S. China, Taiwan.

Chaetocnema (Tlanoma) yiei n. sp.

Oblong-oval. Pitchy black; antennae yellowish brown; legs yellowish to reddish brown with posterior femora much darker.

Head very finely granulate; vertex almost impunctate, except two or three large punctures which are placed on each side close to eyes; inter-antennal space distinct, wide and feebly raised, and not distinctly separated from frons. Antennae filiform; first joint long, robust, club-shaped; second subequal to third in length but distinctly thicker; fourth very slightly longer than third; fourth to tenth subequal to each other in length but gradually thickened toward apical joints; eleventh longer than tenth and pointed at apex. Pronotum transverse; anterior margin slightly rounded anteriorly, anterior corners thickened, showing nearly right angle, posterior corners showing about 120° and each with a large setigerous puncture; lateral margin rounded, widest at basal margin and narrowed anteriorly; posterior margin rounded posteriorly; dorsum rather closely and distinctly punctate by somewhat elongate punctures and their interstices finely granulate. Scutellum hemispheric and its surface finely granulate. Elytra convex, with regularly arranged eleven longitudinal rows of distinct punctures, including a short scutellar row of punctures; interstices of punctate-striae closely impressed with distinct heart-shaped punctures and distinctly costate on lateral area; elytral epipleurae smooth, shining, with a regularly arranged longitudinal row of punctures, alonging to its interior margin.

Length : 1.8mm.

Holotype : Nanshanchi, Nantou Hsien (30. vi. 1965, S. Kimoto leg.).

Paratypes : 11 exs., same as the holotype.

This new species differs from the known form of this genus occurring in Taiwan, in having the punctures of pronotum elongate. Also, this new species resembles *C. longipunctata* Maulik, from Ceylon, but differs in having the punctures of pronotum finer and sparser.

The scientific name of this new species is dedicated to Prof. Shi-tao Yie, National Taiwan University, Taipei.

Distribution : Taiwan.

***Glaucosphaera cyanea* (Duvivier, 1892)**

Fenchihu, Chiayi Hsien (1 ex., 8. vii. 1965, S. Itô leg.).

Distribution : India, Indo-China, Taiwan.

***Pseudoliprus kurosawai* (Nakane, 1958)**

Sungkang (=Taataka), Nantou Hsien (1 ex., 5. v. 1965, 1 ex., 18. v. 1965, T. Shirôzu leg.).

Fenchihu, Chiayi Hsien (1 ex., 13. iv. 1965, T. Shirôzu leg.).

Alishan, Chiayi Hsien (1 ex., 27. vii. 1966, H. Kamiya leg.).

This species is here recorded from Taiwan for the first time.

Distribution : Japan, Taiwan.

***Pseudoliprus saigusai* n. sp.**

Body yellowish to reddish brown, with apical margins of abdominal segments and meso- and metathorax in part blackish in various degrees.

Vertex convex, wrinkled, and with some scattered yellowish hairs, mostly on anterior portion, frontal tubercles indistinct, antennal space very narrow and




Fig. 2. Male genitalia : a, *Pseudoliprus kurosawai* Nakane (Alishan);
b, *P. saigusai* n. sp.

with a deep longitudinal impression at middle. Antennae long, somewhat shorter than length of body; first joints longest, robust, somewhat club-shaped; second shortest, almost $2/3$ as long as first and much slender; third distinctly slender than second and $1\frac{1}{5}$ times as long as second; third to fifth subequal to each other in length and shape; sixth very slightly shorter and slightly robuster than fifth; seventh almost $4/5$ as long as sixth but distinctly robuster; seventh to tenth subequal to each other in length and shape; eleventh $1\frac{1}{5}$ times as long as tenth but its apex pointed. Pronotum subcylindrical, almost as long as wide, feebly constricted slightly behind middle, anterior portion slightly broader than posterior portion; surface convex, with a feebly impressed transverse furrow slightly behind middle, closely covered with yellowish hairs and shallow punctures. Scutellum trigonate, rounded at apex, surface closely covered with setigerous punctures. Elytra elongate, broader than pronotum at base, slightly widened posteriorly; surface convex, closely covered with fine pubescence, and with a broad transverse depression a little before middle, punctures arranging in longitudinal rows large and deep.

Length: 3.0–4.0 mm.

Holotype: Tungpu—Tzuchung, Lulinshan, Nantou Hsien (10. iv. 1965, T. Saigusa leg.).

Paratopotypes: 2 exs., same as the holotype.

Paratypes: Yushankou (=Niitakaguchi) —Alishan (=Arisan), Chiayi Hsien (1 ex., 6. iv. 1967, T. Shirôzu leg.). Alishan, Chiayi Hsien (1 ex., 9. iv. 1965, S. Miyamoto leg.).

This new species somewhat resembles *P. nigrinus* (Jacoby) and *hirtus* (Baly) in having the scutellum covered with the setigerous punctures but distinctly differs in having the body much elongate and larger, and the different characteristics of the male genitalia.

Distribution: Taiwan.

***Lipromorpha shirozui* n. sp.**

Yellowish brown, head blackish brown, pronotum dark brown, scutellum and basal $2/3$ of sutural margin of elytra broadly blackish, meso- and metathorax and abdomen entirely blackish.

Vertex convex, partly wrinkled, and with about twenty distinct setigerous pores of punctures, frontal tubercles vertical, separated to each other by a deep median longitudinal impression. Antennae very slender, long, slightly shorter than length of body; first joint robust, longest, club-shaped, second shortest, about half as long as first; third $1\frac{1}{5}$ times as long as second in length but much slender; fourth subequal to third in length and shape; fifth $1\frac{1}{4}$ times as long as third in length; sixth to tenth gradually widened toward terminal joints; sixth to ninth subequal to each other in length; tenth almost $5/6$ as long as ninth; eleventh $1\frac{1}{2}$ times as long as tenth and its apex pointed. Pronotum subcylindrical, slightly longer than wide, strongly constricted slightly behind middle, anterior portion slightly broader than posterior portion; surface convex, with a deep transverse groove slightly behind middle, closely covered with yellowish hairs and shallow punctures. Scutellum trigonate, rounded at apex,




Fig. 3. Male genitalia: a, *Lipromorpha shirozui* n. sp.; b, *L. montana* (Chûjô) (Fenchihu); c, *L. difficilis* (Chen) (Nanshanchi).

surface finely granulate. Elytra elongate, broader than pronotum at base, slightly widened posteriorly; surface convex, with a broad transverse depression a little before middle, covered with large and deep punctures, punctures arranging in longitudinal rows, each interstices of punctate-striae with a row of yellowish pubescence; each femur incrassate, posterior ones especially so.

Length: 2.0-2.2 mm.

Holotype: Sungkang, Nantou Hsien (29. vi. 1965, T. Shirôzu leg.).

Paratopotypes: 1 ex., same as the holotype.

Paratypes: Sungkang, Nantou Hsien (1 ex., 5. v. 1965, 1 ex., 24. vi. 1965, T. Shirôzu leg.). Sungkang-Tsifeng, Nantou Hsien (2 exs., 29. 1965, S. Kimoto leg.).

This new species closely resembles *L. difficilis* Chen, but differs in having the different coloration of elytra, head, pronotum and ventral surface, and the different characteristics of male aedeagus. Also from *L. montana* Chûjô, in having the interstices of the elytral punctate-striae not costate.

Distribution: Taiwan.

***Lipromorpha montana* Chûjô, 1935**

Peikeng (=Hokukô) —Hushan (=Kaminoshima-Onsen), Miaoli Hsien (1 ex., 11. iv. 1967, T. Shirôzu leg.).

Fenchihu, Chiayi Hsien (1 ex., 10. iv. 1965, 1 ex., 13. iv. 1965, T. Shirôzu leg.; 1 ex., 12. iv. 1965, S. Miyamoto leg.).

5-10 km S. of Fenchihu, Chiayi Hsien (1 ex., 11. iv. 1965, Y. Hirashima leg.).

Distribution: Taiwan.

***Lipromorpha difficilis* (Chen, 1934)**

Neihu, Taipei Hsien (5 exs., 31. iii. 1965, S. Miyamoto leg.).

Nanshanchi, Nantou Hsien (2 exs., 30. iv. 1965, S. Kimoto leg.).

Distribution: China, Taiwan, Ryukyu Is.

***Lipromela formosana* Ohno, 1960**

Hohuanshan, Hualien Hsien (12 exs., 16. vii. 1966, B.-S. Chang leg.).

Huanshan (=Heiganzan), Taichung Hsien—Pinan (=Pianananbu), Ilan Hsien (1 ex., 19. vii. 1932, T. Esaki leg.).

Tsifeng, Nantou Hsien (6 exs., 10. vii. 1966, H. Kamiya leg.).

Sungkang, Nantou Hsien (14 exs., 16. vii. 1966, H. Kamiya leg.).

Fenchihu, Chiayi Hsien (2 exs., 22. vii. 1966, H. Kamiya leg.).

Alishan (=Arisan), Chiayi Hsien (1 ex., 6. vii. 1965, S. Kimoto leg.; 13 exs., 18. viii. 1947, J. L. Gressitt leg.).

Tungpu—Tzuchung, Chiayi Hsien (1 ex., 10. iv. 1965, T. Saigusa leg.).

Distribution: Taiwan.

***Lipromima minuta* (Jacoby, 1885)**

Proc. Zool. Soc. Lond. 1885: 725 (*Liprus*) (Japan)

=*Lipromima fulvipes* Chûjô, 1935, Trans. Nat. Hist. Soc. Formosa 25: 399 (Taiwan: Musha, Kappanzan) (with var. *bicolor* Chûjô). **New Synonymy**

Wulai, Taipei Hsien (1 ex., 4. vii. 1966, H. Kamiya leg.).

Nanshanchi, Nantou Hsien (2 exs., 30. iv. 1965, S. Kimoto leg.).

Distribution: China, Taiwan, Japan.

***Pseudodera xanthospila* Baly, 1862**

Wushe—Yushih, Nantou Hsien (2 exs., 3. v. 1965, T. Shirôzu leg.).

Distribution: Japan, China, Taiwan.

***Xuthea orientalis* Baly, 1865**

Fenchihu, Chiayi Hsien (1 ex., 7. vii. 1965, T. Nakane leg.).

This species is here recorded for the first time from Taiwan.

Distribution: N. India, Burma, SW China, Korea, Taiwan.




Fig. 4. Male genitalia: a, *Micrepitrix coomani* Laboissière (Pahsien-shan); b, *M. okinawana* Kimoto & Gressitt (Okinawa); c, *ibid.* (Kenting).

***Micrepitrix coomani* Laboissière, 1933**

Pahsienshan (=Hassenzan), Taichung Hsien (2 exs., 26. vi. 1934, J. L. Gressitt leg.).

This species is here recorded for the first time from Taiwan.

Distribution: N. Vietnam, Hainan Is., Taiwan.

***Micrepitrix okinawana* Kimoto & Gressitt, 1966**

Kenting, Pingtung Hsien (1 ex., 3. vii. 1965, S. Kimoto leg.).

This species is here recorded for the first time from Taiwan. The aedeagus of this specimen is almost the same as the one illustrated by Kimoto & Gressitt (1966) from Okinawa. However, this specimen differs from the Okinawa specimens in having the dorsum entirely black.

Distribution: Taiwan, Ryukyu Is.

***Sangariola hirashimai* n. sp.**

Head, pronotum and scutellum reddish brown, elytra yellowish brown, with sutural margin broadly and lateral margins narrowly blackish, underside and elytral epipleurae dark reddish brown, antennae entirely pitchy black, legs pitchy black with posterior femora paler.

Body elongate, subparallel-sided. Head with vertex convex, partly granulate, distinctly punctate, with a few yellowish pubescence; frontal tubercles elongate, large, extending to inter-antennal space and separated from vertex by a deep transverse impression, surface rugged and with a few distinct punctures.

Antennae stout, almost $\frac{2}{3}$ as long as length of body, closely furnished with white pubescence; first joint stout, longest, somewhat club-shaped; second shortest, almost half as long as first and much slender; third $1\frac{2}{3}$ times as long as second; fourth $1\frac{1}{5}$ times as long as third; fifth subequal to fourth in length and shape; sixth almost $\frac{5}{6}$ as long as fifth, and sixth to tenth subequal to each other in length but gradually widened toward terminal joints; eleventh almost $1\frac{1}{4}$ times as long as tenth but its apex pointed. Pronotum subquadrate, almost $1\frac{1}{3}$ times as broad as long, anterior margin almost straight, basal margin rounded posteriorly at sides and emarginate at middle, lateral margins rounded, widest at $\frac{1}{3}$ from anterior margin and gradually narrowed anteriorly and posteriorly; surface convex, very sparsely covered with fine yellowish pubescence, and closely impressed with large punctures, and also with five somewhat irregular but well-defined impressions, first one situated near and parallel to basal margin, second one near and parallel




Fig. 5. Male genitalia of *Sangariola hirashimai* n. sp.

to anterior margin, third one longitudinal, and situated at middle and connected with posterior impression, each of last two ones longitudinal, and near and parallel to lateral margin and connected with basal impression. Scutellum trapezoidal, surface wrinkled, sparsely but distinctly covered with setigerous punctures. Elytra elongate, parallel-sided, sparsely covered with fine yellowish pubescence; surface convex, closely impressed with large and deep punctures, which are arranged in eleven regular rows, interstices of any elytral punctate striae not costate.

Length: 3.6-5.0 mm.

Holotype: Alishan, Chiayi Hsien (9. iv. 1965, Y. Hirashima leg.).

Paratype: Alishan, Chiayi Hsien (1 ex., 6. vii. 1965, T. Nakane leg.).

This new species differs from the other known species of this genus in having the interstices of any elytral punctate striae not costate.

Distribution: Taiwan.

***Sangariola fortunei* (Baly, 1888)**

Alishan, Chiayi Hsien (1 ex., 6. vii. 1965, Y. Kurosawa leg.).

Distribution: Taiwan, China, Korea.

***Sangariola punctatostriata* (Motschulsky, 1860)**

= *Sangariola costata* Chûjô, 1935, Trans. Nat. Hist. Soc. Formosa 35: 394 (Taiwan: Musha, Kappanzan). **New Synonymy**

Sungan (=Nihonmatsu)—Peikeng (=Hokukô), Miaoli Hsien (1 ex., 10. iv. 1967, T. Shirôzu leg.; 1 ex., 10. iv. 1967, B.-S. Chang leg.).

Tahu (=Taiko)—Sungan (=Nihonmatsu), Miaoli Hsien (1 ex., 9. iv. 1967, T. Shirôzu leg.).

Distribution: Japan, Korea, N. China, Taiwan? Ryukyu Is.

***Ophrida spectabilis* (Baly, 1862)**

Taipei (=Taihoku), Taipei Hsien (3 exs., 9. viii. 1921, T. Esaki leg.).

Nanshanchi, Nantou Hsien (1 ex., 21. v. 1965, 1 ex., 27. v. 1965, 3 exs., 17. vi. 1965, T. Shirôzu leg.; 1 ex., 1. vi. 1965, B.-S. Chang leg.; 2 exs., 30. vi. 1965, T. Nakane leg.; 1 ex., 30. vi. 1965, S. Kimoto leg.).

Distribution: Taiwan, China.

***Ophrida scaphoides* (Baly, 1865)**

Nanshanchi, Nantou Hsien (1 ex., 17. v. 1965, 1 ex., 21. v. 1965, 1 ex., 30. vi. 1965, T. Shirôzu leg.; 1 ex., 30. vi. 1965, Y.-I. Chu leg.; 1 ex., 17. vi. 1965, 2 exs., 21. vi. 1965, B.-S. Chang leg.; 9 exs., 30. vi. 1965, S. Kimoto leg.; 1 ex., 30. vi. 1965, Y. Kurosawa leg.).

Penpuchi, Nantou Hsien (1 ex., 14. vii. 1965, H. Kamiya leg.).

Chihpen, Taitung Hsien (2 exs., 8. viii. 1965, H. Kamiya leg.).

Distribution: Indo-China, China, Taiwan.

***Podontia lutea* (Olivier, 1790)**

Wulai, Taipei Hsien (1 ex., 4. viii. 1965, Y.-I. Chu leg.).

Taipingshan (=Taiheizan), Ilan Hsien (1 ex., 7. viii. 1934, J. L. Gressitt leg.).

Distribution: Burma, Indo-China, Taiwan.

Nisotra orbiculata (Motschulsky, 1866)

Pinglin, Taipei Hsien (14 exs., 23. iv. 1965, Y. Kurosawa leg.).

Kuangyin, Taoyuan Hsien (12 exs., 10. ix. 1966, B.-S. Chang leg.).

Fenchihu, Chiayi Hsien (1 ex., 12. iv. 1965, T. Saigusa leg.).

Chihpen, Taitung Hsien (4 exs., 10. viii. 1965, H. Kamiya leg.).

Distribution: China, Taiwan, Indo-China, Thailand, Burma, India, Andaman Is., Java, Nicobars.

Neorthaea flavicornis Chen, 1933

Tahu (=Taiko) —Sungan (=Nihonmatsu), Miaoli Hsien (6 exs., 9. iv. 1967, T. Shirôzu leg.).

Pahsienshan (=Hassenzan), Taichung Hsien (1 ex., 26. vi. 1934, J. L. Gressitt leg.).

Puli (=Hori), Nantou Hsien (1 ex., 7. v. 1934, J. L. Gressitt leg.).

Penpuchi, Nantou Hsien (11 exs., 13. vii. 1966, H. Kamiya leg.).

Nanshanchi, Nantou Hsien (1 ex., 26. vi. 1965, T. Shirôzu leg.; 1 ex., 12. vii. 1966, H. Kamiya leg.).

Chiaoliping, Chiayi Hsien (10 exs., 13. iv. 1965, T. Saigusa leg.).

Fenchihu, Chiayi Hsien (14 exs., 10. iv. 1965, 13 exs., 11. iv. 1965, 3 exs., 12. iv. 1965, T. Shirôzu leg.; 53 exs., 10. iv. 1965, Y. Hirashima leg.; 44 exs., 12. iv. 1965, S. Miyamoto leg.; 5 exs., 12. iv. 1965, S. Miyamoto leg.; 44 exs., 12. iv. 1965, T. Saigusa leg.; 5 exs., 10. iv. 1965, M. Sasakawa leg.; 20 exs., 7. vii. 1965, 4 exs., 8. vii. 1965, T. Nakane leg.; 8 exs., 8. vii. 1965, R. Ishikawa leg.; 3 exs., 7. vii. 1965, Y. Kurosawa leg.; 1 ex., 8. vii. 1965, T. Yamasaki leg.; 7 exs., 7. vii. 1965, S. Kimoto leg.; 6 exs., 22. vii. 1966, 9 exs., 23. vii. 1966, 22 exs., 24. vii. 1966, 18 exs., 25. vii. 1966, H. Kamiya leg.).

5-10 km S. of Fenchihu, Chiayi Hsien (3 exs., 11. iv. 1965, Y. Hirashima leg.).

Alishan, Chiayi Hsien (2 exs., 6. vii. 1965, S. Kimoto leg.; 10 exs., 28. vii. 1966, H. Kamiya leg.).

Chulu, Taitung Hsien (12 exs., 11. viii. 1966, H. Kamiya leg.).

Chihpen, Taitung Hsien (2 exs., 8. viii. 1966, 7 exs., 10. viii. 1966, H. Kamiya leg.).

Kuantzuling, Tainan Hsien (1 ex., 7. iv. 1965, T. Saigusa leg.).

Amping, Tainan Hsien (1 ex., 9. ix. 1921, T. Esaki leg.).

Distribution: Taiwan.

Neorthaea nisotroides Chen, 1933

Wulai (=Urai), Taipei Hsien (6 exs., 1. iv. 1932, J. L. Gressitt leg.; 1 ex., 6. vii. 1966, H. Kamiya leg.).

Taipingshan, Ilan Hsien (2 exs., 3. viii. 1967, B.-S. Chang leg.).

Ilan, Ilan Hsien (1 ex., 19. viii. 1966, H. Kamiya leg.).

Tahu (=Taiko) —Sungan (=Nihonmatsu), Miaoli Hsien (2 exs., 9. iv. 1967, T. Shirôzu leg.).

Sungan (=Nihonmatsu) —Peikeng (=Hokukô), Miaoli Hsien (3 exs., 10. iv.

1967, T. Shirôzu leg.).

Sungan, Miaoli Hsien (1 ex., 9. iv. 1967, B.-S. Chang leg.).

Penpuchi, Nantou Hsien (3 exs., 13. vii. 1966, 1 ex., 14. vii. 1966, H. Kamiya leg.).

Meichi, Nantou Hsien (2 exs., 13. v. 1965, 1 ex., 16. v. 1965, T. Shirôzu leg.).

Nanshanchi, Nantou Hsien (1 ex., 27. v. 1965, 4 exs., 20. vi. 1965, 1 ex., 26. vi. 1965, T. Shirôzu leg.; 14 exs., 30. vi. 1965, S. Kimoto leg.; 1 ex., 30. vi. 1965, S. Itô leg.; 1 ex., 30. vi. 1965, T. Nakane leg.; 1 ex., 30. vi. 1965, Y. Kurosawa leg.; 1 ex., 30. vi. 1965, T. Yamasaki leg.; 14 exs., 12. vii. 1966, H. Kamiya leg.).

Wushe—Yushih, Nantou Hsien (3 exs., 3. v. 1965, T. Shirôzu leg.).

Sungkan, Nantou Hsien (7 exs., 5. v. 1965, 2 exs., 18. v. 1965, 5 exs., 10. vi. 1965, 3 exs., 24. vi. 1965, T. Shirôzu leg.; 5 exs., 16. vii. 1966, H. Kamiya leg.).

Sungkan—Tsifeng, Nantou Hsien (12 exs., 29. vi. 1965, S. Kimoto leg.; 3 exs., 29. vi. 1965, T. Nakane leg.; 2 exs., 29. vi. 1965, Y. Kurosawa leg.).

Chihsinliao, Chiayi Hsien (33 exs., 15. iv. 1965, T. Shirôzu leg.; 3 exs., 13. iv. 1965, 2 exs., 15. iv. 1965, T. Saigusa leg.).

Fenchihu, Chiayi Hsien (11 exs., 10. iv. 1965, S. Miyamoto leg.; 1 ex., 10. iv. 1965, T. Shirôzu leg.; 1 ex., 7. vii. 1965, T. Nakane leg.).

Chulu, Taitung Hsien (3 exs., 11. viii. 1966, H. Kamiya leg.).

Kuantzuling, Tainan Hsien (23 exs., 6. iv. 1965, T. Shirôzu leg.; 2 exs., 7. iv. 1965, T. Saigusa leg.).

Kaoshifat (=Kusukusu), Pingtung Hsien (1 ex., 13. iv. 1932, J. L. Gressitt leg.).

Distribution: Taiwan.

***Sinocrepis micans* Chen, 1933**

Yangmingshan, Taipei Hsien (3 exs., 25. v. 1965, K. Morimoto leg.; 18 exs., 24. vi. 1965, 18 exs., 26. vi. 1965, S. Kimoto leg.).

Shuangchi, Hsintien, Taipei Hsien (1 ex., 23. vi. 1965, T. Yamasaki leg.).

Taipei, Taipei Hsien (1 ex., 18. vi. 1965, S. Kimoto leg.).

Neihu, Taipei Hsien (1 ex., 31. iii. 1965, S. Miyamoto leg.).

Puli, Nantou Hsien (1 ex., 8. vii. 1966, 1 ex., 11. vii. 1966, H. Kamiya leg.).

Wushe, Nantou (2 exs., 14. vi. 1961, S. Uéno leg.).

Kuantzuling, Tainan Hsien (1 ex., 6. iv. 1965, T. Saigusa leg.).

Kenting, Pingtung Hsien (3 exs., 2. vii. 1965, 2 exs., 3. vii. 1965, S. Kimoto leg.).

Distribution: China, Taiwan.

***Liprus punctatostriatus* Motschulsky, 1860**

Alishan, Chiayi Hsien (1 ex., 5. vii. 1965, Y. Kurosawa leg.).

This species is here recorded for the first time from Taiwan.

Distribution: Japan, Korea, N. China, Taiwan.

***Hemipyxis plagioderoides flaviabdominalis* Chûjô, 1965**

Wulai (=Urai), Taipei Hsien (1 ex., 1. v. 1932, J. L. Gressitt leg.; 1 ex., 17. iv. 1965, S. Miyamoto leg.; 1 ex., 17. iv. 1965, T. Saigusa leg.).

Taipingshan (=Taiheizan), Ilan Hsien (1 ex., 8. v. 1933, K. Satô leg.).

Tahu (=Taiko) —Sungan (=Nihonmatsu), Miaoli Hsien (2 exs., 9. iv. 1967, T. Shirôzu leg.).

Sungan (=Nihonmatsu) —Peikeng (=Hokukô), Miaoli Hsien (1 ex., 10. iv. 1967, T. Shirôzu leg.).

Penpuchi, Nantou Hsien (1 ex., 2. v. 1965, T. Shirôzu leg.).

Nanshanchi, Nantou Hsien (1 ex., 28. iv. 1965, 6 exs., 30. iv. 1965, T. Shirôzu leg.).

Wushe (=Musha), Nantou Hsien (2 exs., 20. v. 1932, J. L. Gressitt leg.; 2 exs., 14. vi. 1961, S. Uéno leg.; 1 ex., 31. v. 1965, B.-S. Chang leg.).

Wushe—Yushih, Nantou Hsien (2 exs., 3. v. 1965, T. Shirôzu leg.).

Sungkang, Nantou Hsien (3 exs., 5. v. 1965, 3 exs., 18. v. 1965, 3 exs., 19. v. 1965, 2 exs., 31. v. 1965, 1 ex., 1. vi. 1965, 1 ex., 10. vi. 1965, T. Shirôzu leg.; 1 ex., 2. vi. 1965, B.-S. Chang leg.; 6 exs., 29. vi. 1965, T. Yamasaki leg.; 2 exs., 29. vi. 1965, T. Nakane leg.).

Sungkang—Tsifeng, Nantou Hsien (7 exs., 29. vi. 1965, S. Kimoto leg.; 4 exs., 29. vi. 1965, Y. Kurosawa leg.; 1 ex., 29. vi. 1965, S. Itô leg.).

Tsifeng, Nantou Hsien (1 ex., 10. vii. 1966, H. Kamiya leg.).

Meifeng, Nantou Hsien (2 exs., 18. v. 1965, B.-S. Chang leg.).

Fenchihu, Chiayi Hsien (2 exs., 13. iv. 1965, T. Shirôzu leg.; 4 exs., 12. iv. 1965, T. Saigusa leg.; 1 ex., 8. vii. 1965, S. Itô leg.; 1 ex., 9. vii. 1965, Y. Kurosawa leg.).

5-10 km S. of Fenchihu, Chiayi Hsien (21 exs., 11. iv. 1965, Y. Hirashima leg.).

Alishan, Chiayi Hsien (4 exs., 6. vii. 1965, T. Nakane leg.; 3 exs., 6. vii. 1965, S. Kimoto leg.; 1 ex., 6. vii. 1965, Y. Kurosawa leg.).

Distribution: Taiwan.

Hemipyxis balyi (Bates, 1866)

Yangmingshan, Taipei Hsien (5 exs., 25. v. 1965, K. Morimoto leg.; 1 ex., 24. vi. 1965, S. Kimoto leg.).

Wulai, Taipei Hsien (2 exs., 4. vii. 1965, H. Kamiya leg.; 3 exs., 22. viii. 1966, H. Kamiya leg.).

Ilan, Ilan Hsien (1 ex., 16. iv. 1965, Y. Hirashima leg.; 3 exs., 19. viii. 1966, H. Kamiya leg.).

Chaopanshan (=Kappanzan), Taoyuan Hsien (1 ex., 17. ix. 1921, T. Esaki leg.; 2 exs., 17. iv. 1966, B.-S. Chang leg.).

Penpuchi, Nantou Hsien (3 exs., 2. v. 1965, T. Shirôzu leg.).

Nanshanchi, Nantou Hsien (1 ex., 28. iv. 1965, 2 exs., 30. iv. 1965, 4 exs., 21. v. 1965, 2 exs., 8. vi. 1965, 1 ex., 26. vi. 1965, T. Shirôzu leg.; 2 exs., 30. vi. 1965, S. Kimoto leg.; 1 ex., 30. vi. 1965, T. Nakane leg.; 1 ex., 12. vii. 1965, H. Kamiya leg.).

Wushe—Yushih, Nantou Hsien (3 exs., 3. v. 1965, T. Shirôzu leg.).

Sungkang—Tsifeng, Nantou Hsien (1 ex., 29. vi. 1965, S. Kimoto leg.).

Chuchi, Chiayi Hsien (1 ex., 14. iv. 1965, S. Miyamoto leg.).

Fenchihu, Chiayi Hsien (1 ex., 12. iv. 1965, S. Miyamoto leg.; 1 ex., 7. vii. 1965, S. Kimoto leg.; 1 ex., 8. vii. 1965, Y.-I. Chu leg.).

Batakan—Tabito in Tailukohsia (=Taroko), Hualien Hsien (2 exs., 30. vii.

1932, T. Esaki leg.).

Liyuchih, Hualien Hsien (1 ex., 14. viii. 1966, H. Kamiya leg.).

Kueitzuchiao (=Kuraru), Pingtung Hsien (3 exs., 31. viii. 1921, T. Esaki leg.).

Distribution: Taiwan, China.

***Hemipyxis quadripustulata* (Baly, 1876)**

Shuangchi, Hsintien, Taipei Hsien (1 ex., 23. vi. 1965, T. Yamasaki leg.).

Wulai, Taipei Hsien (1 ex., 4. vii. 1966, H. Kamiya leg.).

Ilan, Ilan Hsien (4 exs., 16. iv. 1965, Y. Hirashima leg.).

Sungan, Miaoli Hsien (1 ex., 9. iv. 1967, B.-S. Chang leg.).

Penpuchi, Nantou Hsien (1 ex., 13. vii. 1966, H. Kamiya leg.).

Meichi, Nantou Hsien (3 exs., 13. v. 1965, 1 ex., 16. v. 1965, B.-S. Chang leg.).

Wushe—Yushih, Nantou Hsien (1 ex., 3. v. 1965, T. Shirôzu leg.).

Chiaoliping, Chiayi Hsien (1 ex., 15. iv. 1965, T. Shirôzu leg.; 1 ex., 13. iv. 1965, T. Saigusa leg.).

Fenchihu, Chiayi Hsien (7 exs., 12. iv. 1965, S. Miyamoto leg.; 3 exs., 11. iv. 1965, T. Saigusa leg.; 2 exs., 8. vii. 1965, T. Nakane leg.; 1 ex., 7. vii. 1965, S. Itô leg.; 1 ex., 7. vii. 1965, S. Kimoto leg.).

5–10 km S. of Fenchihu, Chiayi Hsien (2 exs., 11. iv. 1965, M. Sasakawa leg.).

Chihpen, Taitung Hsien (1 ex., 10. viii. 1966, H. Kamiya leg.).

Kuantzuling, Tainan Hsien (3 exs., 6. iv. 1965, T. Saigusa leg.).

Distribution: Taiwan, S. China. Correction: In my previously published paper on the Taiwan Chrysomelidae (1966, Esakia 5: 21–38), the explanation of fig. 5, c, *H. quadrimaculata* (Baly) (page 36) should be read as *H. quadripustulata* (Baly).

***Hemipyxis foveolata* (Chûjô, 1958)**

Yangmingshan, Taipei Hsien (1 ex., 26. vi. 1965, S. Kimoto leg.).




Fig. 6. Male genitalia: a, *Hemipyxis balyi* (Bates) (Chientapu=Zentai); b, *H. foveolata* (Chûjô) (Takao); c, *H. quadripustulata* (Baly) (Chiasien=Kôsempo).

- Taipei (=Taihoku), Taipei Hsien (1 ex., 9. viii. 1921, T. Esaki leg.).
 Ilan, Ilan Hsien (2 exs., 19. viii. 1966, H. Kamiya leg.).
 Nanshanchi, Nantou Hsien (1 ex., 28. iv. 1965, 1 ex., 21. v. 1965, 1 ex., 26. vi. 1965, T. Shirôzu leg.; 1 ex., 30. vi. 1965, S. Kimoto leg.).
 Chuchi, Chiayi Hsien (2 exs., 14. iv. 1965, S. Miyamoto leg.).
 Chihsinliao, Chiayi Hsien (4 exs., 15. iv. 1965, T. Shirôzu leg.; 2 exs., 15. iv. 1965, T. Saigusa leg.).
 Kuantzuling, Tainan Hsien (5 exs., 6. iv. 1965, T. Saigusa leg.; 3 exs., 7. iv. 1965, Y. Hirashima leg.; 2 exs., 6. iv. 1965, S. Miyamoto leg.).
 Chirifu, Kaohsiung Hsien (2 exs., 18. v. 1934, J. L. Gressitt leg.).
 Liyuchih, Hualien Hsien (6 exs., 15. viii. 1966, H. Kamiya leg.).
 Buruburu—Kaimosu, nr. Kuanshan (=Kanzangoe), Taitung Hsien (1 ex., 12. viii. 1932, T. Esaki leg.).
 Chulu, Taitung Hsien (1 ex., 9. viii. 1966, H. Kamiya leg.).
 Kenting, Pingtung Hsien (1 ex., 4. iv. 1965, T. Saigusa leg.).
 Distribution: Taiwan, China, Ryukyu Is.




Fig. 7. Male genitalia: a, *Hemipyxis formosana* (Chûjô) (Nanshanchi); b, *H. quadrimaculata* (Jacoby) (Meiyuan).

***Hemipyxis quadrimaculata* (Jacoby, 1892)**

- Ilan, Ilan Hsien (1 ex., 19. viii. 1966, H. Kamiya leg.).
 Nanshanchi, Nantou Hsien (1 ex., 30. vi. 1965, T. Yamasaki leg.).
 Fenchihu, Chiayi Hsien (1 ex., 7. vii. 1965, 1 ex., 8. vii. 1965, S. Kimoto leg.; 1 ex., 7. vii. 1965, T. Nakane leg.; 1 ex., 8. vii. 1965, R. Ishikawa leg.; 1 ex., 9. vii. 1965, Y. Kurosawa leg.; 1 ex., 8. vii. 1965, T. Yamasaki leg.; 1 ex., 7. vii. 1965, Y.-I. Chu leg.).

Distribution: Sikkim, Burma, China, Taiwan.

***Hemipyxis changi* n. sp.**

Head, pronotum and scutellum shining black, elytra blackish blue; ventral

surface entirely pitchy black; antennae dark brown with four basal joints much paler; legs dark reddish brown.

Vertex wrinkled, nearly impunctate, except for a pair of large foveae situated near interior margins of eyes and a few distinct punctures surrounding large foveae; frontal tubercles convex, subquadrate, contiguous, hardly separated from vertex by a straight, shallow impression; surface smooth, shining. Antennae slender, long, nearly $\frac{2}{3}$ as long as length of body; first joint long, robust, club-shaped; second nearly $\frac{2}{3}$ as long as first and much slender; third as long as second but much slender; fourth $1\frac{1}{5}$ times as long as third in length, fifth subequal to fourth in length and shape, sixth to tenth subequal to each other in length but gradually thickened toward apical joints; eleventh $1\frac{4}{5}$ times as long as tenth and its apex pointed. Pronotum transverse, twice as wide as long, anterior margin almost straight, posterior margin rounded posteriorly, lateral margins widest at base and gradually narrowed toward anterior margin, anterior corners thickened and pointed laterally, basal corners widely rounded, anterior and posterior corners each with a seta-bearing pore, dorsal surface finely and not so closely punctate. Scutellum trigonate, its apex rounded, surface smooth, shining, almost impunctate. Elytra convex, lateral margins rounded, surface closely and distinctly punctate.

Length: 2.0-2.8 mm.

Holotype: Sungan, Miaoli Hsien (9. iv. 1967, B.-S. Chang leg.).

Paratopotypes: 4 exs., same as the holotype.

Paratypes: Tahu (=Taiko) —Sungan (=Nihonmatsu), Miaoli Hsien (4 exs., 9. iv. 1967, T. Shirôzu leg.). Sungkang, Nantou Hsien (1 ex., 31. v. 1965, T. Shirôzu leg.). Tungpu—Tzuchung, Lulinshan, Nantou Hsien (1 ex., 10. iv. 1965, T. Saigusa leg.). Yushankou (=Niitakaguchi) —Alishan (=Arisan), Chiayi Hsien (1 ex., 6. iv. 1967, T. Shirôzu leg.). Ryukyu Is.: Yonehara, Ishigaki Is. (1 ex., 15. iii. 1964, S. Kimoto leg.).

This new species somewhat resembles *H. plagioderoides* (Motschulsky), in having the similar coloration and the general morphological characteristics, but differs in having the length of the body much smaller, vertex nearly impunctate, the surface of frontal tubercles smooth, and the different coloration of antennae and legs.

Distribution: Ryukyu Is. (Ishigaki), Taiwan.

Hemipyxis yasumatsui n. sp.

Head, pronotum, scutellum and ventral surface shining black, elytra yellowish brown, with sutural and lateral margins blackish, but in one specimen elytra entirely brownish; antennae, legs and meso- and metathorax entirely pitchy black; abdomen dark brown.

Vertex convex, smooth, shining, nearly impunctate, except for a pair of large foveae situated near interior margin of eyes; frontal tubercles convex, subquadrate, contiguous, separated from vertex by a straight, shallow impression, surface smooth, shining. Antennae slender, long, nearly $\frac{2}{3}$ as long as length of body; first joint long, robust, club-shaped; second nearly $\frac{2}{3}$ as long as

first and much slender; third $1\frac{1}{5}$ times as long as second and much slender; fourth $1\frac{1}{3}$ times as long as third in length; fifth subequal to fourth in length and shape; sixth $\frac{5}{6}$ as long as fifth and subequal to seventh in length and shape; eighth very slightly shorter than seventh and subequal to ninth in length and shape; ninth subequal to tenth in length and shape; eleventh $1\frac{1}{3}$ times as long as tenth and its apex pointed. Pronotum transverse, almost twice as wide as long, anterior margin rounded posteriorly, lateral margins widest slightly before basal margin and gradually narrowed toward anterior margin, posterior margin widely rounded posteriorly; anterior corners thickened, basal corners widely rounded, anterior and posterior corners each with a seta-bearing pore, dorsal surface impressed with very minute punctures. Scutellum trigonate, its apex rounded, surface smooth, shining, almost impunctate. Elytra convex, lateral margins rounded, surface closely and distinctly punctate.

Length: 5.0 mm.

Holotype: Taipingshan (=Taiheizan), Ilan Hsien (23. x. 1932, K. Satô leg.).

Paratype: Fenchihu, Chiayi Hsien (1 ex., 13. iv. 1965, T. Shirôzu leg.).

This new species resembles *H. flavipennis* Baly, in having the general coloration of the dorsal surface, but differs in having the vertex almost impunctate, the surface of frontal tubercles smooth, and the punctation of pronotum much finer.

Distribution: Taiwan.

Hemipyxis formosana Chûjô, 1936

Taipingshan (=Taiheizan), Ilan Hsien (1 ex., 5. vii. 1934, J. L. Gressitt leg.).

Hsinhsing, Miaoli Hsien (1 ex., 28. vi. 1967, B.-S. Chang leg.).

Nanshanchi, Nantou Hsien (1 ex., 12. vii. 1966, H. Kamiya leg.).

Fenchihu, Chiayi Hsien (1 ex., 8. vii. 1965, T. Nakane leg.; 1 ex., 8. vii. 1965, R. Ishikawa leg.; 1 ex., 8. vii. 1965, T. Yamasaki leg.; 1 ex., 22. vii. 1966, 1 ex., 23. vii. 1966, 2 exs., 25. vii. 1966, H. Kamiya leg.).

Distribution: Taiwan.