

[REVIEW]

Body Plan of Sea Urchin Embryo: An Ancestral Type Animal

Koji Akasaka* and Hiraku Shimada

Laboratory of Molecular Genetics, Department of Mathematical and Life Sciences,
Graduate School of Science, Hiroshima University,
Higashi-Hiroshima 739-8526, Japan

ABSTRACT—Sea urchin embryos are thought to possess a body plan characteristic of early deuterostomes. Sea urchins contain homologs of *Otx*, *Lim*, *T-brain* and *Hox* gene cluster, which are involved in head and segment formation in vertebrate development, although the sea urchin has not evolved a head or segments. We described here that sea urchin *Otx* is involved in various aspects of early development and that the *Hox* genes do not obey spatial colinearity in sea urchin embryo. The *Otx* and *Hox* genes seem to be used subsequently for head formation and determining the anteroposterior axis respectively during chordate evolution. We propose that the Precambrian was a period where these regulatory genes were utilized in many different combinations during animal development, leading to the evolution of a wide range of body plans, many of which were successful. We also discuss the role of chromatin boundaries and the mechanism of cell specification along animal vegetal axis, especially differentiation of the large micromere progeny, which are the prospective primary mesenchyme cells and play a role as an organizer in sea urchin embryos.

INTRODUCTION

The animals alive today that are known to science are composed of approximately one million species displaying a tremendous diversity. All the basic types of multicellular animals burst on the scene in an evolutionary frenzy called the Cambrian explosion about 540 million years ago. Genes involved in morphogenesis are well conserved and the function of most of the genes are shared among all animal phyla, thus the divergent animals are thought to be derived from a common ancestor. It has been suggested that the early animals had already obtained almost all the genes responsible for the morphogenesis by the Precambrian period and that the evolution of genetic regulatory programs produced divergent animals. The molecular mechanisms involved in giving rise to the diverse animal phyla alive today needs to be elucidated, however the origin of animals has been elusive because of the dearth of animal fossils below the Precambrian-Cambrian boundary. In 1998, animals considerably older than the Cambrian finally began being found in the fossil record from the 570 million-year old Doushantuo phosphorites in southern China, and they were preserved in a way that

reveals details down to the cellular level (Xiao *et al.*, 1998; Li *et al.*, 1998). The key to the exquisite preservation is calcium phosphate which is known for its faithful replication of delicate tissues. The Doushantuo phosphorite contained large populations of animal embryos in cleavage stages as well as algae and sponges. The manner of cleavage of the embryo suggested that the Doushantuo fossils are most probably bilaterian. Although gastrulae or later developmental stages have not yet been identified, these findings suggested that they could be broadly equivalent to blastulae or planuloids *sensu* Haeckel (1874), embryos of microscopic animals as envisioned by Davidson *et al.* (1995) and that the morphology of the Precambrian animals resemble larvae of marine invertebrates alive today.

We have been studying molecular mechanisms of early development using sea urchin embryos which are thought to possess a body plan characteristic of early deuterostomes (Davidson *et al.*, 1995). Sea urchins are simple deuterostomes with bilaterally symmetrical, enterocoelous larvae. *Otx* and *Hox* genes, which are well conserved in metazoans, are involved in the head formation and anteroposterior positional information of segments in chordates respectively. How were these transcription factors used in the ancestral animals which had not yet evolved head and segments? In this review, we focus on the function of *Otx* and *Hox* genes in sea urchin early development, and also we describe the factors involved in the

* Corresponding author: Tel. +81-824-24-7447;
FAX. +81-824-24-7327.
E-mail: koji@ipc.hiroshima-u.ac.jp

primary mesenchyme cell differentiation and new findings using an advantage of sea urchin embryos as an animal model.

Development of sea urchin embryo

In sea urchin embryogenesis it has been suggested that the initial territories are specified by a combination of asymmetric distribution of cytoplasmic determinants and cell-cell interactions. During cleavage, a maternally regulated, reproducible pattern of cell divisions partitions the egg cytoplasm among blastomeres that consequently have defined sizes and orientations relative to each other. The geometric precision of cleavage restricts the range of cell-cell interactions that take place in the normal embryo, with the result that the fates of blastomeres at different positions along the animal-vegetal axis are reproducible and predictable, although most blastomeres have the potential to assume a wide variety of fates until early gastrula stage. At the 16-cell stage, animal-vegetal polarity is morphologically evident. From animal to vegetal are arrayed tiers of eight mesomeres, four macromeres, and four micromeres. At the 32-cell stage, the mesomeres have divided to give two tiers of 8 cells, while in the vegetal hemisphere the macromeres have divided to give one tier of 8 cells, and the micromeres have divided to give large and small micromeres (Fig. 1). At the 60-cell stage blastomeres clonally originated from founder cells divide the embryo into five distinct territories: small micromeres, large micromeres, vegetal plate, oral ectoderm, and aboral ectoderm (Fig. 2). The vegetal plate develops into archenteron and secondary mesen-

chyme cells which produce muscle and pigment cells in later stage. The territories are identified by the expression of specific marker genes and their cell lineages (Davidson, 1989, 1991) (Fig. 2).

The large micromeres are thought to play a role as an organizer and initiate a cascade of signal transduction toward overlying cells (Davidson, 1989). The large micromeres induce the overlying macromere progenies, specifying the vegetal plate (Ransick and Davidson, 1993, 1995). A complete respecification of cell fate occurs when 16-cell stage micromeres from the vegetal pole of a donor embryo are implanted into the animal pole of an intact recipient embryo (Fig. 1). Thus, the large micromeres, which are the prospective primary mesenchyme cells (PMCs) play a key role in the cell fate specification and axis determination during sea urchin embryogenesis.

The mesomeres at 16-cell stage display a character similar to the animal cap of amphibian embryo. The mesomeres are normally fated to form epithelial cells. When the mesomeres are combined with micromeres and/or large micromeres, the chimera embryo develops three germ layers and forms an almost normal embryo (Livingston and Wilt, 1990; Khaner and Wilt, 1991) (Fig. 1).

Macromeres at 16-cell stage develop into ectoderm, endoderm and secondary mesenchyme cells (SMCs). Borders between ectoderm and endoderm and between endoderm and SMCs are negotiated by inductive interactions among macromere progeny during the late cleavage-to-mesenchyme




Fig. 1. Development of sea urchin embryo. Micromere descendant cells (gray) play a key role in sea urchin development. Micromeres (black) implanted into animal pole induce secondary gut.




Fig. 2. Territories and territory specific markers of the sea urchin embryo. Ars; arylsulfatase (Akasaka *et al.*, 1990; Mitsunaga-Nakatsubo *et al.*, 1998), Spec2a; calmodulin like protein (Cox *et al.*, 1986), Cyllla; cytoplasmic actin (Hardin *et al.*, 1988), EctoV; hyaline layer glycoprotein protein (Coffman and McClay, 1990), Hpoe; hyaline layer glycoprotein (Yoshikawa, 1997), Endo16; cell adhesion protein (Nocente-McGrath *et al.*, 1989), SM30; spicule matrix protein (George *et al.*, 1991; Akasaka *et al.*, 1994), SM50; spicule matrix protein (Benson *et al.*, 1987; Katoh-Fukui *et al.*, 1992), msp130; lipid-anchored glycoprotein (Parr *et al.*, 1990), HpTa; Brachyury (Harada *et al.*, 1995).

blastula period (Angerer and Angerer, 2000).

During the cleavage stage, the embryonic cells form an epithelial ball with monolayer-cells (blastula). After hatching, the cells around the vegetal pole migrate into blastocoel (mesenchyme blastula). These cells are PMCs which later form spicules. Then the cells around vegetal pole begin to invaginate to form the archenteron (gastrula). The invagination site represents the anus. At the late gastrula stage, SMCs appear at the tip of the archenteron. The SMCs develop into muscles and pigment cells later. The archenteron is bent and opens into a stomodeum. The oral side of the ectoderm is referred to as oral ectoderm, and the other side is referred to as aboral ectoderm (prism).

Hox7 and Hox11/13b are involved in pattern formation in sea urchin embryos

It is well known that *Hox* gene complexes have highly conserved function in determining the anteroposterior axis. Sea urchins also possess a single *Hox* gene complex containing 10 genes (Martinez *et al.*, 1999) (Fig. 3a). However, in the embryo, only two *Hox* genes (*Hox7* and *Hox11/13b*) are expressed (Angerer *et al.*, 1989; Dobias *et al.*, 1996). The expression of *Hox7* protein is restricted to the aboral ectoderm, and *Hox11/13b* expression is restricted to oral ectoderm, endoderm and SMCs in sea urchin embryos after the gastrula stage (Fig. 3b).




Fig. 3. Sea Urchin Hox gene complex. (a) Structure of sea urchin Hox gene complex. (b) Expression territories of *Hox7* and *Hox11/13b* in sea urchin embryo. (c) Ectopic expression changes the morphology and expression pattern of marker genes.

The nuclear localization of Hox7 and Hox11/13b was demonstrated by immunostaining (Dobias *et al.*, 1996; Ishii *et al.*, 1999), supporting the idea that both Hox proteins are involved in transcriptional control. With the aim of gaining insight into the role of Hox7 and Hox11/13b in the sea urchin development, we performed Hox7 and Hox11/13b overexpression experiments by injecting mRNAs of Hox7 and Hox11/13b. The overexpression of Hox7 represses the development of oral ectoderm, endoderm and mesenchyme cells. On the contrary, overexpression of Hox11/13b represses the development of aboral ectoderm and PMCs. The data suggests that Hox7 and Hox11/13b are expressed in distinct non overlapping territories, and over expression of either one inhibits territory specific gene expression in the domain of the other (Fig. 3c).

Since the Hox7 overexpressing embryos develop into epithelial balls consisting of aboral ectoderm cells alone, it is likely that Hox7 is part of the circuit responsible for the differentiation of aboral ectoderm cells. However, it has been shown that the aboral ectoderm cells have initiated a tissue-specific program of gene expression before the Hox7 message achieves a significant fraction of its peak abundance (Angerer L. M. *et al.*, 1989). Thus, Hox7 can not be involved in the initial specification of aboral ectoderm. Quantitative RT-PCR reveals that the overexpression of Hox7 does not activate aboral-ectoderm specific *Ars* gene expression. Thus, it does not seem likely that Hox7 is involved in aboral ectoderm differentiation by activating aboral ectoderm-specific genes.

Nickel (Sunderman, 1989) ions have been implicated as chemical reagents which disturb the signal transduction. It has been shown that the intracellular signaling mediated by calcium ions is involved in the differentiation of oral-aboral ectoderms (Akasaka *et al.*, 1997). It is likely that cell to cell signals regulate aspects of the ectodermal patterning during early development. The Hox7 overexpressing embryo is resistant to oral ectoderm induction by NiCl_2 . Thus, Hox7 seems to be involved in the last crucial part of the cascade which results in the oral-aboral differentiation. In Hox11/13b overexpressing embryos, both oral ectoderm (EctoV) and endoderm specific (*Endo16*) genes are expressed in the epithelial cells, however the morphological differentiation of archenteron and oral ectoderm was not observed. It seems likely that factors other than Hox11/13b are necessary for the complete development of endoderm and oral ectoderm.

We propose that an important function of both of Hox7 and Hox11/13b genes in sea urchin embryo is to maintain specific territorial gene expression by each one, and their function does not depend on cell position along the axis of the embryo (Ishii *et al.*, 1999). Deschamps and Wijgerde (1993) reported that there are two phase of Hox gene expression in mouse development. One phase of Hox gene expression, which is well known, takes place at the early somite stage. Another phase of Hox gene expression takes place at the gastrula stage, much earlier than the somite stage. They showed that Hox-2.3 and Hox-2.4 start to express at the late streak stage at the allantois and the most posterior part of the

streak. In *C. elegans* also, the Hox genes are reported to be responsible for the specification of cell fate in a position independent manner (Cowing and Kenyon, 1996; Wittmann *et al.*, 1997).

Recently Arenas-Mena *et al.* (2000) have shown a spatially sequential and colinear arrangement of expression domains of five posterior genes of the Hox cluster in the somatocoel during larval stage. The sequence of Hox gene expression patterns within the somatocoel was cross orientated to the adult anterior-posterior axis. They also reported that a remarkable number of apparent co-options of Hox gene use in developing structure. The findings in sea urchin development, and those from work in *C. elegans* and mouse early embryo suggest that Hox gene cluster may originally have had functions in the establishment of the territories or cell-fate, and do not necessarily obey spatial colinearity. In some embryos, the genes are used subsequently for determining the anteroposterior axis at the somite stage.

Otx is involved in various aspects in sea urchin development

The Otx gene family is a member of the bicoid class homeobox genes. The product contains a well conserved homeodomain which has a lysine at position 51 that confers DNA-binding specificity for the sequence motif TAATCC/T (Hanes and Brent, 1989, 1991; Treisman *et al.*, 1989). Orthodenticle (*Otd*) cDNA was first isolated from *Drosophila* as a homeobox protein related to head formation (Finkelstein *et al.*, 1990; Cohen and Jurgens, 1990; Finkelstein and Perrimon, 1990). Subsequently, cDNA homolog (*Otx*) of *Otd* were isolated from various vertebrates. These Otx proteins also have been shown to be essential for patterning the anteriormost aspects of the brain in vertebrates. The mouse has two types of Otx which are expressed in restricted regions of developing brain and are responsible for head formation (Simeone *et al.*, 1993; Matsuo *et al.*, 1995). In zebrafish three types of Otx are present in the developing diencephalon and midbrain (Mori *et al.*, 1994). In sea urchin development, two isoforms of Otx are expressed (Mao *et al.*, 1994; Sakamoto *et al.*, 1997). The first type of Otx protein, referred to as an




Fig. 4. Structure of Otx gene and protein. (a) Structure of Otx protein. HD: homeodomain, DBD: DNA binding domain, AD: activation domain. (b) Structure of Otx gene. Boxes represent exons. Gray regions in the box represent coding regions. Lower panel indicates splicing pattern of Otx gene products.




Fig. 5. Spatial expression pattern of *HpArs*, *HpOtxE* and *HpOtxL*. Regions of the embryo where genes are expressed are indicated by the dark gray.

early-type Otx (*HpOtxE*), appears in early development and gradually decreases in amount by the gastrula stage. The second type of Otx protein, referred to as a late-type of Otx (*HpOtxL*), appears at the blastula stage and remains until gastrula stage. The nucleotide sequence reveals that the homeobox and downstream regions through to the C-terminus are identical in the two types of *HpOtx* proteins, while the N-terminal region has different polypeptides (Sakamoto *et al.*, 1997) (Fig. 4a). These distinct Otx proteins are generated from a single gene by altering the transcription start site and by an alternative splicing (Kiyama *et al.*, 1998) (Fig. 4b). Whole-mount in situ hybridization using isoform specific probes reveals a complex and dynamic change of expression patterns in the three germ layers (Li *et al.*, 1997; Mitsunaga *et al.*, 1998) (Fig. 5), suggesting that the Otx is not merely required for the differentiation of specific territories. *Otx* genes seem to be involved in the various aspect of sea urchin early development.

The overexpression of Otx isoforms alters the fate of entire embryonic cells to ectoderm cells (Mao *et al.*, 1996; Mitsunaga *et al.*, 1998). Furthermore, disrupting Otx function by dominant repression causes a specific inhibition of aboral ectoderm- and endoderm-specific gene expression and blocks the formation of aboral ectoderm and endoderm cell types (Li *et al.*, 1999). These dramatic changes of morphogenesis produced by overexpression and reduced function indicate that Otx plays an important role in the sea urchin embryos. We suggest that *Otx* may originally have had functions in various aspects in early development, and the gene was used subsequently for head formation during chordate evolution.

Otx target gene

Despite important role of Otx in embryogenesis, the direct target genes have not been demonstrated in any other species other than sea urchins. *HpArs* (Sakamoto *et al.*, 1997; Kiyama *et al.*, 1998), *Spec2a* (Gan *et al.*, 1995) and *Endo16* (Yuh *et al.*, 1998) have been reported as direct target genes of Otx in sea urchin embryo. *HpArs* is transcriptionally activated late in blastula, and after the gastrula stage it is expressed exclusively in the aboral ectoderm throughout embryonic development (Sasaki *et al.*, 1988; Akasaka *et al.*,

1990a, 1990b). In order to detect cis-regulatory elements responsible for the transcriptional activity of genes, we introduce reporter(*Luc*)-fusion constructs into fertilized eggs. To analyze the function of transcription factors *in vivo*, we introduce effector(transcription factor)-constructs together with reporter-constructs (Kiyama *et al.*, 1998). Microinjection is a very useful technique to introduce DNA into fertilized egg; however, it does have some disadvantages: a certain amount of skill is required to perform the microinjection, and the number of embryos that can be injected at one session is limited to approximately 1000. We applied the particle gun method to sea urchin embryos and obtained excellent expression of the introduced DNA (Akasaka *et al.*, 1995). We introduce DNA into approximately 100,000 eggs per one shot. The expression of the construct of interest can be normalized by concomitantly-introducing a reference construct; CMV *Renilla* luciferase, which allows highly quantitative experiment. When the linearized plasmid DNAs are introduced into fertilized egg, they form random concatamers, and during the early development of the embryos replicate repeatedly (McMahon *et al.*, 1985). The stable incorporation of exogenous DNA into the chromosomal DNA of embryos occurs, and the exogenous DNAs persist until postmetamorphosis juvenile stage (Flytzanis *et al.*, 1985).

In gene transfer experiments, a 229-bp fragment, referred to as C15, in the first intron of *HpArs* was found to have enhancer elements (Iuchi *et al.*, 1995). This region contains a tandem repeat of core consensus sequences of orthodenticle-related protein (Otx) binding sites, which serve as the major source of positive control for *HpArs* gene (Sakamoto *et al.*, 1997) (Fig. 6a). The time course of expression of *HpOtxL* is similar to that of *Ars* gene. Thus, *HpOtxL* is very likely to be involved in the activation of the *Ars* gene. In order to eliminate the contribution of the endogenous *HpOtx* to the expression of the reporter, we used a Gal4 DNA binding domain-*HpOtx* fusion protein together with a reporter containing Gal4 binding sites and lacking *HpOtx* sites. This should provide a test of the ability of *HpOtx* to transactivate the *Ars* promoter-enhancer without the background contribution by the endogenous *HpOtx*. The reporter construct was generated by ligating Gal4




Fig. 6. Structure of HpArs enhancer and transactivation assay with the Gal4-Otx fusion protein. (a) Structure of HpArs enhancer. Sequences in open boxes indicate Otx binding sites. Sequences in gray boxes indicate CTF binding sites. (b) Upper panel indicates the structure of the expression construct of the Gal4-Otx fusion protein. Lower panel indicates luciferase reporter gene driven by different combination of cis-elements, and the expression levels.

binding site (UAS), (in stead of *Ars* enhancer C15 which contains Otx sites) and an *Ars* promoter, (*Ars194*) in which the Otx binding site has been deleted, conjoined to the luciferase reporter gene (Fig. 6b). The different construct combinations were introduced into fertilized eggs, which were cultured to the gastrula stage. HpOtxL successfully activates the *Ars* promoter by 4 fold, but that HpOtxE does not under the same conditions. The Gal4-HpOtxL fusion protein does not activate the expression of the luciferase reporter construct in which the Gal4 binding sites are deleted. The results indicate that Gal4-HpOtxL fusion protein activates *Ars* promoter-directed gene activity at the UAS site.

Otx activates *ars* gene with CAAT binding protein

We thought the 4 fold expression activated by Gal4-HpOtxL fusion protein was much less than endogenous *Ars* enhancer C15 activity. We wished to know if regions of the C15 enhancer other than Otx sites are needed for maximum enhancer activity. Therefore, we generated an enhancer in which UAS was substituted for the Otx sites, and in which 5' and/or 3' deletions ($\Delta 5'$ and/or $\Delta 3'$) were made in the modified C15 enhancer. When both of the 5' and 3' regions of the C15 enhancer were fused to UAS (C15(UAS)), the fusion enhancer C15(UAS) shows approximately 3-fold greater activity than UAS alone without the expression of Gal4-HpOtxL fusion protein. Furthermore, the activity of the *Ars* promoter(*Ars194*) bearing C15(UAS) was enhanced 5 fold by the expression of Gal4-HpOtxL fusion protein (Fig. 6b). The results indicate that

both the 5' region and 3' region of the C15, in addition to Otx sites, are required for a maximum response to HpOtxL.

Besides the Otx sites, the enhancer C15 contains two CAAT sequences (Fig. 6a). We sought to determine the functional properties of these two sites. In order to analyze the role of Otx sites and CAAT sequences precisely, site-directed mutagenesis was employed to mutate these sequences. When one of the CAAT sequences was changed but the Otx sites were unaltered, a 15–34% reduction in the activity was observed. A significant (60%) decrease in the activity was observed when both CAAT sequences were mutated but the Otx sites were kept intact. When both Otx sites and CAAT sequences were mutated, the activity was reduced to background, indicating that together, both CAAT sequences and Otx sites are essential for the maximal enhancer activity. Furthermore, we have demonstrated that nuclear proteins prepared from embryos at mesenchyme blastula stage bind to the CAAT sequences in the enhancer C15 (Kiyama *et al.*, 2000).

Functional domains in Otx

Because the fusion enhancer C15(UAS) responds appropriately to the expression of Gal4-HpOtxL fusion protein, we used C15(UAS) as a target sequence to address the question of which domains of HpOtxL are responsible for the activation of the *Ars* promoter. We fused portions of HpOtxL with the Gal4-DNA-binding domain and analyzed the ability of these fusion proteins to activate a luciferase reporter gene driven by C15(UAS)-*Ars194* in sea urchin embryos. The de-

letion analysis revealed that C-terminal region between amino acid 218–238 of HpOtxL are required for the transactivation, and that the N-terminal region of HpOtxL is responsible for the enhancement of transactivation of the *Ars* promoter. The N-terminal region of the other isoform, HpOtxE, represses the activity of its C-terminal region (Fig. 4a). These data suggest that Otx regulates the transcription of a target *Ars* gene by working cooperatively with a CAAT sequence binding factor(s), and that the transactivation domain and the regulation domain exists in the C terminal region and the N terminal region respectively.

Specificity of Otx target genes

The enhancer region of *Spec2a*, which is also specifically expressed in the aboral ectoderm, contains two Otx sites and two CAAT sequences (Gan and Klein, 1993). Considered together with the findings in the present study, it is possible that both Otx and CAAT sites are responsible for the specific expression of genes in the aboral ectoderm (Fig. 6a). On the other hand, the combination of a synthetic Otx site and the basal promoter of *Endo16* suffices to generate endoderm-specific expression (Yuh *et al.*, 1998). Thus Otx regulates different genes in different territories in the embryo, and it is unlikely Otx is responsible for the territory-specific expression of the target genes. A previous report suggested that different transcription factors are involved in the expression of one gene and the factors form a regulatory network to achieve a complex gene expression in development (Yuh *et al.*, 1998). We have suggested that HpOtxL activates *Ars* expression co-operatively with CAAT binding protein. It is likely that co-factor(s) interacting with Otx direct the specificity of the target genes.

Genes involved in animal-vegetal axis patterning

The cell fates of cells in the early embryo are progressively specified along the animal-vegetal axis. Embryos may

inherit positional information in the form of asymmetrically distributed maternal determinants that are positioned during oogenesis. When unfertilized eggs are bisected through the equator and the two halves are fertilized, the animal half gives rise to an incompletely differentiated epithelial ball, while the vegetal half can produce a normal pluteus larva with derivatives of endoderm, mesoderm, and ectoderm (Horstadius, 1939; Maruyama *et al.*, 1985). This experiment suggests the presence of determinants of fate in the vegetal pole and the requirement for cell-cell interactions to complete the fate specification of animal blastomeres. Recently, members of Wnt signaling system, such as GSK-3 β , β -catenin and Tcf/Lef, have been reported to be factors which are involved in pattern formation of sea urchins along the animal-vegetal axis. Tcf/Lef is a partner of β -catenin in transcriptional regulation (Clevers and van de Wetering, 1997). Introduction into sea urchin embryos of a dominant negative form of Tcf/Lef that lacks the β -catenin binding domain produces animalized phenotypes (Huang *et al.*, 2000; Vonica *et al.*, 1999). The converse is also true: a constitutively active Tcf consisting of the strong transcription activation domain of VP16, vegetalized embryos (Vonica *et al.*, 1999). Activation of GSK-3 β promotes the degradation of β -catenin through a phosphorylation, ubiquitination and proteosome pathway, resulting in low level of free cytoplasmic β -catenin. Overexpression of GSK-3 β causes severe animalization. Again the converse is true: expression of dominant negative (kinase-dead) GSK-3 β blocks the turnover, leading to over accumulation of β -catenin in nuclei and vegetalization (Emily-Fenouil *et al.*, 1998) (Fig. 7a). Treatment of embryos with LiCl, an inhibitor of GSK 3 β , expands the domain of nuclear β -catenin into the presumptive ectoderm territory (Logan *et al.*, 1999) and the embryos are vegetalized. The dominant negative Tcf/Lef can reverse the vegetalizing effect of LiCl (Vonica *et al.*, 1999). An animal-vegetal asymmetry in the levels of nuclear β -catenin is seen in the embryos after 16 or 32-cell stage (Logan *et al.*, 1999) (Fig. 7b). The




Fig. 7. Animal vegetal axis patterning. (a) Factors involved in animal vegetal axis patterning. Li; Li ion, kd-GSK3 β ; GSK3 β lacking kinase domain, dn-TCF/Lef; TCF/Lef lacking β -catenin binding domain. A; animal pole, V; vegetal pole. (b) Animal-vegetal asymmetry in the levels of nuclear β -catenin. A; animal pole, V; vegetal pole.

expression of *Wnt8* is also first detected after 16 or 32-cell stage (Wikramanayake personal communication). However, no asymmetrical distribution of nuclear β -catenin is observed before the 8 or 16-cell stage, thus mechanisms other than the conventional Wnt-signaling pathway must be present in order to establish the animal-vegetal axis in the egg. Maternally inherited RNAs and/or proteins could be responsible for the formation of animal-vegetal axis in the egg.

Transcription factors involved in primary mesenchyme cell differentiation

Micromeres, which are located at the vegetal pole in 16-cell stage embryos, divide at the 5th cleavage into 4 large micromeres and 4 small micromeres. Descendants of the large micromeres immigrate into the blastocoel and give rise to PMCs, which are responsible for skeletogenesis in the pluteus larvae (reviewed by Davidson *et al.*, 1998), and play a key role in the cell fate specification and axis determination during sea urchin embryogenesis. Previous data suggested that the large micromeres are autonomously specified to become PMCs by maternally inherited determinants (Okazaki, 1975; Kitajima and Okazaki, 1980). An important question in sea urchins embryogenesis is the identity and function of the proposed maternal determinants. To gain a foothold for the elucidation of the cascade of micromere specification initiated by maternally inherited factors, we have isolated transcription factors responsible for PMC-specification. We found that *HpEts*, which encodes the Ets-related transcription factor of sea urchin, *Hemicentrotus pulcherrimus*, play an important role in the differentiation of PMCs (Kurokawa *et al.*, 1999). *HpEts* is expressed exclusively in the PMCs after the blastula stage (Fig. 8a). The overexpression of *HpEts* by injecting the mRNA into fertilized eggs results in the alteration of the fate of all the embryonic cells into migrating cells, which is characteristic of PMCs, and the migrating cells expressed spicule matrix protein SM50 (Fig. 8b). Also the animal cap (mesomere) assay revealed that isolated mesomeres, which is fated to form ectoderm in the normal embryo, are transformed into skeletogenic PMCs when the animal cap is derived from eggs injected with *HpEts* mRNA (Fig. 8c). On the other hand, embryos injected with a truncated *HpEts* mRNA, which encodes the ets-DNA binding domain lacking the N terminal region, develop normally except for the skeletogenic PMCs. Spicule formation was repressed in these embryos (Fig. 8d). Quantitative reverse transcribed polymerase chain reaction used for monitoring the expression of various cell type specific marker genes revealed that the embryos injected with *HpEts* mRNA showed a low level of *HpArs* (aboral ectoderm specific) and *HpEndo16* mRNA (endoderm specific) in comparison to uninjected embryos. On the other hand, the level of primary mesenchyme specific *HpSM50* mRNA increased significantly in comparison to that in uninjected embryos. Embryos injected with the truncated *HpEts*-mRNA showed a normal level of *HpArs* (aboral ectoderm specific) and *HpEndo16* mRNA (endoderm specific), but the level of PMC specific *HpSM50* mRNA decreased significantly in comparison to uninjected embryos

(Fig. 8). We also demonstrated that the ets binding site located in the upstream region of SM50 (Kurokawa *et al.*, 1999) and SM30 (Akasaka *et al.*, 1994, Kurokawa *et al.*, 1999, Yamasu and Wilt, 1999) is a positive transcriptional element. These findings suggested that the *HpEts* gene is involved in sea urchin PMC differentiation. However, *HpEts* does not seem to be involved in the organizer activity of primary mesenchyme cells, since the embryos expressing dominant negative *HpEts* form archenteron and SMCs. Recently, we have isolated a sea urchin homolog of *T-brain*, a transcription factor containing T-domain. We referred the gene as *HpTb*. *HpTb* is exclusively expressed in the PMCs. When the expression of *HpTb* is repressed by injecting with anti-sense morpholino oligo, formation of archenteron and the SMCs are not observed in the injected embryo (submitted). We suggest that *HpTb* is involved in the organizer activity of PMCs.

Notch signaling is involved in SMC induction by PMC

It is well known that the micromeres have the capacity to induce a second axis if transplanted to the animal pole, and the absence of micromeres at the vegetal pole results in the failure of macromere progeny to specify SMCs. This suggests that micromeres have the capacity to induce SMCs (Fig. 1).

The evolutionarily conserved Notch intercellular signaling pathway has been shown to play an essential role in the segregation of a diverse array of cell types in both invertebrate and vertebrate embryos (Kimble and Simpson, 1997). The sea urchin homolog of the Notch receptor displays dynamic patterns of expression within both the presumptive SMCs and endoderm during the blastula stage, the time at which these two cell types are thought to be differentially specified (Sherwood and McClay, 1997). The activation of Notch by injecting eggs with mRNA encoding a constitutively active form of Notch results in an increase in SMC specification, while loss or reduction of Notch signaling by overexpression of dominant negative forms of Notch eliminates or significantly decreases SMC specification (Sherwood and McClay, 1999). Transplantation studies show that much of the vegetal hemisphere is competent to receive the induction signal. They suggested that the micromeres induce SMCs, most likely through direct contact with macromere progeny, or at most with cells only a cell diameter away (McClay *et al.*, 2000). The induction is quantitative in that more SMCs are induced by four micromeres than by one. Furthermore, they demonstrated that micromeres requires nuclear β -catenin to exhibit SMC induction activity (Sherwood and McClay, 1999). They suggest that the macromeres first must transport β -catenin to their nuclei in order to be receptive to the micromere inductive signal, and as one consequence the Notch pathway becomes competent to receive the micromere induction signal, and to transduce that signal (Fig. 9). Temporal studies showed that the induction signal is passed from the micromeres to macromere progeny between the eighth and tenth cleavage. This is consistent with the observation previously reported by Minokawa and Amemiya (1999).

Recently, sea urchin *delta* homolog is found to be ex-




Fig. 8. Function of HpEts in sea urchin development. (a) HpEts expressing cells are indicated by the dark gray. Signals from ectoderm are thought to be required for the formation of spicule. (b) Ectopic expression of HpEts results in the alteration of all the embryonic cell-fate into migrating PMCs. The transformed cells produce spicules in vitro in the presence of serum. (c) A chimera composed of one mesomere isolated from an embryo injected with HpEts mRNA, which is indicated with dark gray, and eight mesomeres isolated from a normal embryo. The mesomere progeny injected with HpEts mRNA form spicules. Animal cap (mesomeres) is fated to form ectoderm. (d) Dominant negative expression of HpEts which lacks the activation domain results in a normal prism embryo except for the formation of skeletogenic PMCs. The cells which express the truncated form of HpEts are indicated by light gray.

pressed exclusively in PMCs when the PMCs have the capacities to induce SMCs (Ettenshorn, personal communication). *delta* is known to be a ligand for Notch (Morrison *et al.*, 2000; Bally *et al.*, 2000). It is possible that *delta* is a SMC inducing factor emanated from PMCs.

Boundary of chromatin

A great number of genes are arranged on a single DNA molecule. Since enhancer elements are able to activate promoters at great distances and in an orientation independent manner (Atchison, 1988), the enhancer elements in one gene could influence the promoter of other genes. The chromatin




Fig. 9. Model for endoderm and SMC induction by micromere progeny. The intensity of color in the nucleus indicates the concentration of nuclear β -catenin.

fibers of the eukaryotic genome are thought to be organized into discrete domains. Boundaries may serve to confine enhancers and silencers to act only within a domain, as well as to insulate the genes within a domain from the cis-regulatory elements or chromatin structure of an adjacent domain. A boundary designated as an insulator is defined as a DNA fragment which blocks the activity of enhancer when it is located between the enhancer and the promoter of the gene (Kellum and Schedl, 1992; Chung *et al.*, 1993), and which confers position independent transcription on transgenes stably integrated into the chromosome (Kellum and Schedl, 1991; Kalos and Fournier, 1995) (Fig. 10a).

We have been studying the mechanism of expression of *Ars* gene, which is expressed in a tightly regulated manner (Sasaki *et al.* 1987, Sasaki *et al.*, 1988; Akasaka *et al.*, 1990a, b) during sea urchin development. The *Ars* gene is actively transcribed during post gastrula stages, and the accumulation of arylsulfatase mRNA may reach about 1% of total mRNAs (Sasaki *et al.*, 1988); the C15 enhancer in the first intron is apparently very strong. Therefore, the *Ars* enhancer could influence the promoter of the neighboring genes. We assumed that a boundary of chromatin is located in the flanking region of such an actively transcribed *Ars* gene. In order to find putative insulator elements in the upstream region of *Ars*, we used the well-characterized *Ars* enhancer C15 (225 bps), which contains Otx binding sites as major positive elements (Iuchi *et al.*, 1995; Sakamoto *et al.*, 1997; Kiyama *et al.*, 1998), together with *Ars* promoter (–252 bp to +38 bp) (Akasaka *et al.*, 1994; Morokuma *et al.*, 1997) to drive a luciferase reporter gene. The constructs were introduced in the fertilized eggs by

particle gun and the transcriptional activity was determined at gastrula stage when the *Ars* gene is actively transcribed. We found that a fragment spanning from –2686 bp to –2113 bp of *Ars* gene acts as an insulator (Fig. 10b). Gel mobility shift assay revealed that at least two different proteins extracted from sea urchin embryos bind to the *Ars* insulator sequence.

To determine whether the *Ars* insulator isolated from sea urchin genome functions in other species when coupled with other regulatory elements, we tried the sea urchin insulator in *Drosophila* embryo by employing P element-mediated transformation. We introduced a plasmid construct containing the *even-skipped* stripe enhancer-promoter (Cai and Levine, 1995) with or without the insulator fragment. When the insulator fragment was placed between the enhancer of stripe 3 and stripe 2, the insulator fragment blocked stripe 3 expression from the *eve/lacZ* promoter, but allowed the stripe 2 enhancer to direct the expression of its promoter (Fig. 10c). The simultaneous use of two different stripe enhancers (*eve* stripes 2 and 3) indicate that the enhancer lying distal to the sea urchin insulator is selectively blocked in *Drosophila*.

When a gene is inserted close to a heterochromatic region in a gene transfer experiment, the transgene is subjected to repression by the chromatin environment. It is therefore of interest to determine if the insulator fragment insulates transgenes from repressing chromosomal position effect. We stably transfected HeLa cells with the constructs in which the neomycin-resistant gene driven by a SV40 enhancer-promoter is flanked by the insulator fragments (Fig. 10d). The number of geneticin-resistant colonies increased significantly when the neomycin resistant gene is flanked on both sides by the insu-




Fig. 10. Ars insulator. (a) Definition of insulator. (b) An insulator in the flanking region of the Ars gene. (c) The Ars insulator functions in *Drosophila* embryo. Upper panel indicates the expression pattern of the reporter gene. St.2; stripe 2, St.3; stripe 3. Lower panel indicates the structure of reporter constructs which are introduced into *Drosophila* embryos. St.2; stripe 2 enhancer, St.3; stripe 3 enhancer, INS; Ars insulator. (d) Inhibition of position effect by the Ars insulator. Structure of constructs which are introduced into HeLa cells and the expression. SV40-E-P; enhancer and promoter derived from SV40, Neo^r; neomycin resistant gene, INS; Ars insulator. The activity is represented as number of colonies/plate.

lator fragment. The findings suggest that the sea urchin Ars insulator could overcome the position-dependent repression of transgene expression in mammalian cells. Recently, we have demonstrated that the sea urchin Ars insulator functions in mouse embryo (Takada *et al.*, 2000) and in plant cells (Nagaya *et al.*, 2001). Currently, different boundary sequences have been reported (Kellum and Schedl, 1991; Chung *et al.*, 1993; Michel, *et al.*, 1993; Cai *et al.*, 1995; Kalos and Fournier, 1995; Hagstrom *et al.*, 1996; Zhong *et al.*, 1997; Akasaka *et al.*, 1999). However, no sequence similarity was observed among them. Some of the insulators, such as chicken β -globin insulator (Chung *et al.*, 1993) and the sea urchin Ars insulator reported here, function in species other than those from which they originate. Given these findings, we expect that the mechanism(s) of action of insulators have been conserved throughout evolution, and that the nuclear proteins which bind

to the boundary sequences may recognize higher order DNA structures rather than specific DNA sequences.

Concluding remarks

It has been suggested that by Precambrian period early animals had already obtained almost all the genes responsible for the morphogenesis of animals alive today and that the evolution of different genetic regulatory programs is what produced divergent animals. Phylogenetic studies have revealed that systems involved in morphogenesis are well conserved among divergent animals. However, in some genes, there seem to be certain variety in their usages. Sea urchin possesses homologs of *Otx*, *Lim*, *T-brain* and the *Hox* gene cluster, which are involved in head and segment formation in vertebrate development, although the sea urchin has not evolved a head or segments. We propose that the Precam-

brian was a period where these regulatory genes were utilized in many different combinations during animal development, leading to the evolution of a wide range of body plans, many of which were successful. We described here that sea urchin *Otx* is involved in various aspects of early development and that the Hox genes do not obey spatial colinearity in sea urchin embryo. The *Otx* and *Hox* genes seems to be used subsequently for head formation and determining the antero-posterior axis respectively during chordate evolution. The acquisition of an anterior-posterior axis of the body, especially a head at the most anterior end of the body, seems to be extremely advantageous for survived over a long evolution history. Thus only animals, for instance, arthropod and vertebrate, which used these genes to establish anterior-posterior axis of the body could increase their population and give rise to new species. The animals which did not use these genes for the establishment of the anterior-posterior axis, such as sea urchin, thus could not increase their number. On the other hand, the chromatin boundary, which is a basic mechanism responsible for the regulation of gene expression, seems to be conserved among all the variety of animals. The elucidation of the mechanisms of evolution of body plans is one of the most important problem we face. In order to understand the whole story of evolution of body plans, we need to investigate the functions of many different genes involved in morphogenesis in different marine invertebrates, which are thought of as the ancestral type of animals.

ACKNOWLEDGEMENTS

We thank the Editor-in-Chief of this journal, Professor Norio Suzuki, for recommending us to write this review. The authors wish to express their thanks to members of our laboratory, especially Kazuko Takata, Keiko Mitsunaga-Nakatsubo, Mamoru Ishii, Daisuke Kurokawa, Naoaki Sakamoto, Takae Kiyama. The authors also thank Dr. Brian Livingston (University of Missouri, Kansas City) for his advice in preparation and critical reading of the manuscript, and Dr. H. Katow (Asamushi Marine Biological Station) for supplying live sea urchins. A part of this work was carried out in the Center for Gene Research of Hiroshima University. We thank the Cryogenic Center, Hiroshima University for supplying cryogenic liquid nitrogen. This work was supported in part by Grants-in-Aid for Scientific Research (C2) (No. 11680724) and for Scientific Research on Priority Areas (No. 11152227) to K.A., and for Scientific Research (B) (No. 07458195) to H.S. from the Ministry of Education, Science, Sports and Culture, Japan, and Program for Promotion of Basic Research Activities for Innovative Biosciences.

REFERENCES

- Akasaka K, Akimoto Y, Sato M, Hirano H, Shimada H (1990b) Histochemical detection of arylsulfatase activity in sea urchin embryos. *Dev Growth Differ* 32: 293–298
- Akasaka K, Frudakis TN, Killian CE, George NC, Yamasu K, Khaner O, Wilt FH (1994) Genomic organization of a gene encoding the spicule matrix protein SM30 in the sea urchin *Strongylocentrotus purpuratus*. *J Biol Chem* 269, 20592–20598
- Akasaka K, Nishimura A, Hijikata K, Iuchi Y, Morokuma J, Takahashi M, Shimada H (1995) Introduction of DNA into sea urchin eggs by particle gun. *Mol Mar Biol Biotechnol* 4: 255–261
- Akasaka K, Nishimura A, Takata K, Mitsunaga K, Mibuka F, Ueda H, Hirose S, Tsutsui K, Shimada H (1999) Upstream Element of the Sea Urchin Arylsulfatase Gene Serves as an Insulator. *Cell Mol Biol* 45: 555–565
- Akasaka K, Sakamoto N, Yamamoto T, Morokuma J, Fujikawa N, Takata K, Eguchi S, Shimada H (1994) Corrected structure of the 5' flanking region of arylsulfatase gene of the sea urchin, *Hemicentrotus pulcherrimus*. *Dev Growth Differ* 36: 633–636
- Akasaka K, Ueda T, Higashinakagawa, T, Yamada K, Shimada H (1990a) Spatial pattern of arylsulfatase mRNA expression in sea urchin embryo. *Dev Growth Differ* 32: 9–13
- Akasaka K, Uemoto H, Wilt F, Mitsunaga KM, Shimada H (1997) Oral-aboral ectoderm differentiation of the sea urchin embryos is disrupted in response to calcium ionophore. *Dev Growth Differ* 39: 373–379
- Angerer LM, Angerer RC (2000) Animal-vegetal axis patterning mechanisms in the early sea urchin embryo. *Dev Biol* 218: 1–12
- Angerer LM, Dolecki GJ, Gagnon ML, Lum R, Wang G, Yang Q, Humphreys T, Angerer RC (1989) Progressively restricted expression of a homeo box gene within the aboral ectoderm of developing sea urchin embryos. *Genes Dev* 3: 370–383
- Arenas-Mena C, Cameron A, Davidson EH (2000) Spatial expression of Hox cluster genes in the ontogeny of a sea urchin. *Development* 127: 4631–4643
- Atchison ML (1988) Enhancers: mechanism of action and cell specificity. *Annu Rev Cell Biol* 4: 127–153
- Bally-Cuif L, Goutel C, Wassef M, Wurst W, Rosa F (2000) Coregulation of anterior and posterior mesendodermal development by a hairy-related transcriptional repressor. *Genes Dev* 14: 1664–1677
- Cai H, Levine M (1995) Modulation of enhancer-promoter interactions by insulators in the *Drosophila* embryo. *Nature* 376: 462–463
- Chung JH, Whiteley M, Felsenfeld G (1993) A 5' element of the chicken β -globin domain serves as an insulator in human erythroid cells and protects against position effect in *Drosophila*. *Cell* 74: 505–514
- Cohen SM, Jurgens G (1990) Mediation of *Drosophila* head development by gap-like segmentation genes. *Nature* 346: 482–485
- Cowing D, Kenyon C (1996) Correct *Hox* gene expression established independently of position in *Caenorhabditis elegans*. *Nature* 382: 353–356
- Davidson EH (1989) Lineage-specific gene expression and the regulative capacities of the sea urchin embryo: A proposed mechanism. *Development* 105: 421–445
- Davidson EH (1991) Spatial mechanisms of gene regulation in metazoan embryos. *Development* 113: 1–26
- Davidson EH, Peterson KJ, Cameron RA (1995) Origin of bilaterian body plans; Evolution of developmental regulatory mechanisms. *Science* 270: 1319–1325
- Deschamps J, Wijgerde M (1993) Two phases in the establishment of HOX expression domains. *Dev Biol* 156: 473–480
- Dobias SL, Zhao AZ, Tan H, Bell JR, Maxson R (1996) *SpHbox7*, a new *Abd-B* class homeobox gene from the sea urchin *Strongylocentrotus purpuratus*: Insights into the evolution of Hox gene expression and function. *Dev Dyn* 207: 450–460
- Emily-Fenouil F, Ghiglione C, Lhomond G, Lepage T, Gache C (1998) GSK3 β /shaggy mediates patterning along the animal-vegetal axis of the sea urchin embryo. *Development* 125 : 2489–2498
- Finkelstein R, Smouse D, Capaci TM, Sparadling AC, Perremon N (1990) The *orthodenticle* gene encodes a novel homeo domain protein involved in the development of *Drosophila* nervous system and ocellar visual structures. *Genes Dev* 4: 1516–1527
- Finkelstein R, Perrimon N (1990) The *orthodenticle* gene is regulated by *bicoid* and *torso* and specifies *Drosophila* head development. *Nature* 346: 485–488
- Flytzanis CN, McMahon AP, Hough-Evans BR, Katula-KS, Britten RJ, Davidson EH (1985) Persistence and integration of cloned DNA

- in postembryonic sea urchins. *Dev Biol* 108: 431–442
- Davidson EH, Cameron RA, Ransick A (1998) Specification of cell fate in the sea urchin embryo: summary and some proposed mechanisms. *Development* 125: 3269–3290
- Gan L, Klein WH (1993) A positive cis-regulatory element with a bicoid target site lies within the sea urchin *Spec2a* enhancer. *Dev Biol* 157: 119–132
- Gan L, Mao C, Wikramanayake A, Angerer LM, Angerer RC, Klein WH (1995) An orthodenticle-related protein from *Strongylocentrotus purpuratus*. *Dev Biol* 167: 517–528
- Haeckel E (1874) The gastraea-theory, the phylogenetic classification of the animal kingdom and the homology of the germ-lamel-lae. *Quart J Microsc Sci* 14: 142–165
- Hagstrom K, Muller M, Schedl P (1996) *Fab-7* functions as a chromatin domain boundary to ensure proper segment specification by the *Drosophila* bithorax complex. *Genes Dev* 10: 3202–3215
- Hanes SD, Brent R (1989) DNA specificity of the bicoid activator protein is determined by homeodomain recognition helix residue 9. *Cell* 57: 1275–1283
- Hanes SD, Brent R (1991) A genetic model for interaction of the homeodomain recognition helix with DNA. *Science* 251: 426–430
- Horstadius S (1973) "Experimental Embryology of Echinoderms." Clarendon Press, Oxford
- Ishii M, Mitsunaga-Nakatsubo K, Kitajima T, Kusunoki S, Shimada H, Akasaka K (1999) Hbox1 and Hbox7 are involved in pattern formation in sea urchin embryos. *Dev Growth Differ* 41: 241–252
- Iuchi Y, Morokuma J, Akasaka K, Shimada H (1995) Detection and characterization of the cis-element in the first intron of the *Ars* gene in the sea urchin. *Dev Growth Differ* 37: 373–378
- Kalos M, Fournier RE (1995) Position-independent transgene expression mediated by boundary elements from the apolipoprotein B chromatin domain. *Mol Cell Biol* 15: 198–207
- Kellum R, Schedl P (1991) A position-effect assay for boundaries of higher order chromosomal domains. *Cell* 64: 941–950
- Kellum R, Schedl P (1992) A group of *scs* elements function as domain boundaries in an enhancer-blocking assay. *Mol Cell Biol* 12: 2424–2431
- Kimble J, Simpson P (1997) The LIN-12/Notch signaling pathway and its regulation. *Annu Rev Cell Dev Biol* 13: 333–361
- Kiyama T, Sasai K, Takata K, Mitsunaga-Nakatsubo K, Shimada H, Akasaka K (2000) CAAT sites are required for the activation of the *H. pulcherrimus* *Ars* gene by Otx. *Dev Genes Evol* 210: 583–590
- Kitajima T, Okazaki K (1980) Spicule formation in vitro by the descendants of precocious micromere formed at the 8-cell stage of sea urchin embryo. *Dev Growth Differ* 22: 265–279
- Khaner O, Wilt F (1991) Interactions of different vegetal cells with mesomeres during early stages of sea urchin development. *Development* 112: 881–890
- Kurokawa D, Kitajima T, Mitsunaga-Nakatsubo K, Amemiya S, Shimada H, Akasaka K (1999) *HpEts*, an ets-related transcription factor implicated in primary mesenchyme cell differentiation in the sea urchin embryo. *Mech Dev* 80: 41–52
- Li CW, Chen JY, Hua TE (1998) Precambrian sponges with cellular structures. *Science* 279: 879–882
- Li X, Chuang C-K, Mao C-A, Angerer LM, Klein WH (1997) Two Otx proteins generated from multiple transcripts of a single gene in *Strongylocentrotus purpuratus*. *Dev Biol* 187: 253–266
- Li X, Wikramanayake AH, Klein WH (1999) Requirement of SpOtx in cell fate decisions in the sea urchin embryo and possible role as a mediator of β -catenin signaling. *Dev Biol* 212: 425–439
- Livingston BT, Wilt FH (1990) Range and stability of cell fate determination in isolated sea urchin blastomeres. *Development* 108: 403–410
- Logan CY, Mille JR, Ferkowicz MJ, McClay DR (1999) Nuclear β -catenin is required to specify vegetal cell fates in the sea urchin embryo. *Development* 126: 345–357
- Mao C, Gan L, Klein WH (1994) Multiple Otx binding sites required for expression of the *Strongylocentrotus purpuratus* *Spec2a* gene. *Dev Biol* 165: 229–242
- Mao CA, Wikramanayake AH, Gan L, Chuang CK, Summers RG, Klein WH. (1996) Altering cell fates in sea urchin embryos by overexpressing SpOtx, an orthodenticle-related protein. *Development* 122: 1489–1498
- Martinez P, Rast JP, Arenas-Mena C, Davidson EH (1999) Organization of an echinoderm Hox gene cluster. *Proc Natl Acad Sci USA* 96: 1469–1474
- Maruyama YK (2000) A sea cucumber homolog of the mouse *T-brain-1* is expressed in the invaginated cells of the early gastrula in *Holothuria leucospilota*. *Zool. Sci.* 17: 383–387
- Matsuo I, Kuratani S, Kimura C, Takada N, Aizawa S (1995) Mouse Otx2 functions in the forebrain and patterning of rostral head. *Genes Dev* 9: 2646–2658
- McClay DR, Peterson RE, Range RC, Winter-Vann AM, Ferkowicz MJ (2000) A micromere induction signal is activated by β -catenin and acts through Notch to initiate specification of secondary mesenchyme cells in the sea urchin embryo. *Development* 127: 5113–5122
- McMahon AP, Flytzanis CN, Hough-Evans BR, Katula KS, Britten RJ, Davidson EH (1985) Introduction of cloned DNA into sea urchin egg cytoplasm: replication and persistence during embryogenesis. *Dev Biol* 108: 420–430
- Michel D, Chatelain G, Herault Y, Harper F, Brun G (1993) H-DNA can act as a transcriptional insulator. *Cell Mol Biol Res* 39: 131–140
- Minokawa T, Amemiya S (1999) Timing of the potential of micromere descendants in echinoid embryos to induce endoderm differentiation of mesomere-descendants. *Dev Growth Differ* 41: 535–547
- Mitsunaga KN, Akasaka K, Sakamoto N, Takata K, Matsumura Y, Kitajima T, Kusunoki S, Shimada H (1998) Differential expressions of sea urchin Otx isoform (HpOtxE and HpOtxL) mRNAs during early development. *Int J Dev Biol* 42: 645–451
- Mori H, Miyazaki Y, Morita T, Nitta H, Mishima M (1994) Different spatio-temporal expressions of these *otx* homeoprotein transcripts during zebrafish embryogenesis. *Mol Brain Res* 27: 2221–231
- Morokuma J, Akasaka K, Mitsunaga KN, Shimada H (1997) A cis-regulatory element within the 5' flanking region of arylsulfatase gene of sea urchin, *Hemicentrotus pulcherrimus*. *Dev Growth Differ* 39: 469–476
- Morrison SJ, Perez SE, Qiao Z, Verdi JM, Hicks C, Weinmaster G, Anderson DJ (2000) Transient Notch activation initiates an irreversible switch from neurogenesis to gliogenesis by neural crest stem cells. *Cell* 101: 499–510
- Nagaya S, Yoshida K, Kato K, Akasaka K, Shinmyo A (2001) Insulator of sea urchin suppresses variation of transgene expression in cultured tobacco cells. *Mol Gen Genet* 265: 405–413
- Okazaki K (1975) Spicule formation by isolated micromeres of the sea urchin embryo. *Am Zool* 15: 567–581
- Ransick A, Davidson EH (1995) Micromeres are required for normal vegetal plate specification in sea urchin embryos. *Development* 121: 3215–3222
- Ransick A, Davidson EH (1993) A complete second gut induced by transplanted micromeres in the sea urchin embryo. *Science* 259: 1134–1138
- Sakamoto N, Akasaka K, Nakatsubo KM, Takata K, Nishitani T, Shimada H (1997) Two isoforms of orthodenticle-related proteins (HpOtx) binds to the enhancer element of sea urchin arylsulfatase gene. *Dev Biol* 181: 284–295
- Sasaki H, Akasaka K, Shimada H, Shiroya T (1987) Developmental timing of synthesis and translation of arylsulfatase mRNA in sea urchin embryo. *Dev Growth Differ* 29: 317–322
- Sasaki H, Yamada K, Akasaka K, Kawasaki H, Suzuki K, Saito A,

- Sato M, Shimada H (1988). cDNA cloning, nucleotide sequence and expression of the gene for arylsulfatase in the sea urchin (*Hemicentrotus pulcherrimus*) embryo. *Eur J Biochem* 177: 9–13
- Sherwood DR, McClay DR (1997) Identification and localization of a sea urchin Notch homologue: insights into vegetal plate regionalization and Notch receptor regulation. *Development* 124: 3363–3374
- Sherwood DR, McClay DR (1999) LvNotch signaling mediates secondary mesenchyme specification in the sea urchin embryo. *Development* 126: 1703–1713
- Simeone A, Acampora D, Mallamaci A, Stornaiuolo A, D'Apice MR, Nigro V, Boncinelli E (1993) A vertebrate gene related to orthodenticle contains a homeodomain of the bicoid class and demarcates anterior neuroectoderm in the gastrulating mouse embryo. *EMBO J* 12: 2735–2747
- Sunderman FW (1989) Mechanisms of nickel carcinogenesis. *Scan J Work Environ Health* 15: 1–12
- Takada T, Iida K, Akasaka K, Yasue H, Torii R, Tsujimoto G, Taira M, Kimura H (2000) Evaluation of heterologous insulator function with regard to chromosomal position effect in the mouse blastocyst and fetus *Mol Reprod Dev* 57: 1–6
- Treisman J, Gonczy P, Vashishtha M, Harris E, Desplan C (1989) A single amino acid can determine the DNA binding specificity of homeodomain proteins. *Cell* 59: 553–562
- Wittmann C, Bossinger O, Goldstein B, Fleischmann M, Kohler R, Brunschwig K, Tobler H, Müller F (1997) The expression of the *C. elegans labial*-like *Hox* gene *ceh-13* during early embryogenesis relies on cell fate and on anteroposterior cell polarity. *Development* 124: 4193–4200
- Xiao S, Zhang Y, Knoll AH (1998) Three-dimensional preservation of algae and animal embryos in a Neoproterozoic phosphorite. *Nature* 391: 553–558
- Yamasu K, Wilt FH (1999) Functional organization of DNA elements regulating *SM30α*, a spicule matrix gene of sea urchin embryos. *Dev Growth Differ* 41: 81–91
- Yuh CH, Bolouri H, Davidson EH (1998) Genomic cis-regulatory logic: experimental and computational analysis of a sea urchin gene. *Science* 279: 1896–1902
- Zhong XP, Krangel M (1997) An enhancer-blocking element between alpha and delta gene segments within the human T cell receptor alpha/delta locus. *Proc Natl Acad Sci USA* 94: 5219–5224

(Received March 9, 2001 / Invited Review)